arXiv:1201.6128v1 [math.LO] 30 Jan 2012

HOD in Natural Models of AD^+

Nam Trang
Department of Mathematics
University of California
Berkeley, California, 94720 USA
http://math.berkeley.edu/∼namtrang
namtrang@math.berkeley.edu.

January 31, 2012

Abstract

The goal of this paper is to compute the full HOD of natural models of AD^+ below "$AD_{\mathbb{R}} + \Theta$ is regular". As part of this computation, we completed the computation of HOD|Θ left open in [3] for Θ a successor in the Solovay sequence.

1 Introduction

We start with the definition of Woodin's theory of AD^+.

Definition 1.1. *AD^+ is the theory $ZF + AD + DC_{\mathbb{R}}$ and*

1. *for every set of reals A, there are a set of ordinals S and a formula φ such that $x \in A \Leftrightarrow L[S, x] \vDash \varphi[S, x]$;*

2. *for every $\lambda < \Theta^1$, for every continuous $\pi : \lambda^\omega \to \omega^\omega$, for every $A \subseteq \mathbb{R}^2$, the set $\pi^{-1}[A]$ is determined.*

Definition 1.2. *Assume AD^+. The Solovay sequence is the sequence $\langle \theta_\alpha \mid \alpha \leq \Omega \rangle$ where*

1. *θ_0 is the sup of ordinals β such that there is an OD surjection from $\mathbb{R}$ onto β;*

---

[1] Θ is the sup of ordinals α such that there is a surjection $\pi : \mathbb{R} \to \alpha$.

[2] We identify $\mathbb{R}$ with ω^ω in this paper.

2. *if $\alpha > 0$ is limit, then $\theta_\alpha = \sup\{\theta_\beta \mid \beta < \alpha\}$;*

3. *if $\alpha = \beta + 1$ and $\theta_\beta < \Theta$, fixing a set $A \subseteq \mathbb{R}$ of Wadge rank θ_β, θ_α is the sup of ordinals γ such that there is an $OD(A)$ surjection from $\mathbb{R}$ onto γ.*

AD^+ is arguably the right theory for models of the form $V = L(\mathcal{P}(\mathbb{R}))$. So in this paper, we assume $V = L(\mathcal{P}(\mathbb{R})) + AD^+ +$ no AD^+ models M containing $\mathbb{R} \cup Ord$ satisfying "$AD_{\mathbb{R}} + \Theta$ is regular"[3]. We call this assumption $(*)$. Our smallness assumption is made because of the fact that for our computation, we'll rely heavily on the theory of hod mice, which is developed in [3] for models satisfying the assumption. This paper computes full HOD in these models.

To put this work in a proper context, we recall a bit of history on the computation of HOD. In $L(\mathbb{R})$ under AD[4], Harrington and Kechris show that HOD $\vDash CH$. Let $\kappa = \omega_1^{L(\mathbb{R})}$. Solovay shows that HOD $\vDash \kappa$ is measurable and Becker shows κ is the least measurable in HOD. These were shown using descriptive set theory. Then Steel in [6] or [9] using inner model theory shows V_Θ^{HOD} is a fine-structural mouse, which in particular implies $V_\Theta^{HOD} \vDash GCH$. Woodin (see [8]) then shows HOD $\vDash GCH + \Theta$ is Woodin and furthermore shows that the full HOD of $L(\mathbb{R})$ is a hybrid mouse where some information about a certain iteration strategy is fed into the model. A crucial point for Steel's and Woodin's computation of HOD is that if $L(\mathbb{R}) \vDash AD$ then $L(\mathbb{R}) \vDash MC$[5]. It's natural to ask whether analogous results hold in the context of $AD^+ + V = L(\mathcal{P}(\mathbb{R}))$ as the HOD computation is an integral part of the structural analysis of AD^+ models and plays an important role in the core model induction. Woodin has shown that under this assumption HOD $\vDash CH$. Recently, Grigor Sargsyan in [3], assuming $(*)$, proves Strong Mouse Capturing (SMC) and computes V_Θ^{HOD} for Θ being limit in the Solovay sequence and $V_{\theta_\alpha}^{HOD}$ for $\Theta = \theta_{\alpha+1}$ in a similar sense as above under the assumption $(*)$.

This paper extends Sargsyan's work to the computation of full HOD under $(*)$. There are two main cases. We show that if Θ is θ_0 or is a successor in the Solovay sequence, HOD is of the form $L[\mathcal{M}_\infty][\Sigma_\infty]$, where $\mathcal{M}_\infty$ is a fine structural premouse (hybrid premouse if $\Theta > \theta_0$) extending HOD$|\Theta$. The definition of $\mathcal{M}_\infty$ will be spelled out in detail during the course of the paper. Σ_∞ is a fragment of the strategy for $\mathcal{M}_\infty|\Theta$ on (finte stacks of) normal trees in $\mathcal{M}_\infty$. For clarity, we devote the entire Section 2 to the computation of HOD for AD^+ models satisfying $\Theta = \theta_0$. In Section 3, we bring in the machinery of hod mice developed in

---

[3] Under our hypothesis, "Strong Mouse Capturing" (SMC) holds. This technical notion will be introduced in Sections 2 and 3

[4] It's known that if $L(\mathbb{R}) \vDash AD$ then $L(\mathbb{R}) \vDash AD^+$.

[5] MC stands for Mouse Capturing, which is the statement that if $x, y \in \mathbb{R}$, then $x \in OD(y) \Leftrightarrow x$ is in a mouse over y.

[3] and combine it with techniques of Section 2 to compute HOD for AD^+ models satisfying $\Theta = \theta_{\alpha+1}$ for some α. The case Θ is a limit in the Solovay sequence, i.e. $\Theta = \theta_\alpha$ for some limit α, is dealt with in Section 4. There we'll see that the HOD computation is split into two cases depending on whether or not $HOD \vDash cof(\Theta)$ is measurable.

This work is done under the supervision of the author's adviser, Professor John Steel. The author would like to thank him for suggesting this topic, his patience, and numerous helpful advice during the course of this project. The extent to which this paper is in debt to Grigor Sargsyan's work on hod mice will be apparent in Chapters 3 and 4. The author would also like to thank him for numerous suggestions and corrections on an older version of this paper.

2 The $\Theta = \theta_0$ case

2.1 Definitions and Notations

We recall some basic notions from descriptive inner model theory. All the notions and notations used in this section are standard. See [4] and [5] for full details. Here, we'll try to stay as close notationally as possible to those used in [4] and [5]. The reader who are familiar with basic descriptive inner model theory can skip ahead to the actual computation and come back to this when necessary.

Suppose $A \subseteq \mathbb{R}$ and (N, Σ) is such that $N \vDash ZFC - Replacement$ and Σ is an (ω_1, ω_1)-iteration strategy or just ω_1-iteration strategy for N. Suppose that δ is countable in V but is an uncountable cardinal of N and suppose that $T, U \in N$ are trees on δ. We say (T, U) *locally Suslin captures* A over N if for any $\alpha < \delta$ and for N-generic $g \subseteq Coll(\omega, \alpha)$,

$$A \cap N[g] = p[T]^{N[g]} = \mathbb{R}^{N[g]} \backslash p[U]^{N[g]}.$$

We also say that N locally captures A at δ. We say that N locally captures A if N locally captures A at any uncountable cardinal of N. We say (N, Σ) *Sulsin captures* A at δ, or (N, δ, Σ) *Sulsin captures* A, if there are trees $T, U \in N$ on δ such that whenever $i : N \to M$ comes from an iteration via Σ, $(i(T), i(U))$ locally Suslin captures A over M at $i(\delta)$. In this case we also say that $(N, \delta, \Sigma, T, U)$ Suslin captures A. We say $(\mathcal{N}, \Sigma)$ Suslin captures A if for every countable δ which is an uncountable cardinal of N, (N, Σ) Suslin captures A at δ. When δ is Woodin in N, one can perform genericity iterations on N to make various objects generic over an iterate of N. This is where the concept of Suslin capturing becomes interesting and useful. We'll exploit this fact many times in this paper.

Definition 2.1. *Let* $0 \leq k < \omega$ *and* Γ *be a good pointclass* [6]. *A premouse* $\mathcal{N}$ *is* k*-suitable with respect to* Γ *iff there is a strictly increasing sequence* $\langle \delta_i \mid i \leq k \rangle$ *such that*

1. *for all* δ, $\mathcal{N} \vDash$ *"*δ *is Woodin" iff* $\delta = \delta_i$ *for some* $i < 1 + k$*;*
2. $ORD^{\mathcal{N}} = sup(\{(\delta_k^{+n})^{\mathcal{N}} | n < \omega\})$*;*
3. $Lp^{\Gamma}(\mathcal{N}|\xi) \trianglelefteq \mathcal{N}$ *for all cutpoints* ξ *of* $\mathcal{N}$ *where* $Lp^{\Gamma}(\mathcal{N}|\xi) = \cup\{\mathcal{M} \mid \mathcal{N}|\xi \trianglelefteq \mathcal{M} \wedge \rho(\mathcal{M}) = \xi \wedge \mathcal{M}$ *has iteration strategy in* $\Gamma\}$*;*
4. *if* $\xi \in ORD \cap \mathcal{N}$ *and* $\xi \neq \delta_i$ *for all* i*, then* $Lp^{\Gamma}(\mathcal{N}|\xi) \vDash$ *"*ξ *is not Woodin."*

Definition 2.2. *Let* $\mathcal{N}$ *be as above and* $A \subseteq \mathbb{R}$*. Then* $\tau^{\mathcal{N}}_{A,\nu}$ *is the unique standard term* $\sigma \in \mathcal{N}$ *such that* $\sigma^g = A \cap \mathcal{N}[g]$ *for all* g *generic over* $\mathcal{N}$ *for* $Col(\omega, \nu)$*, if such a term exists. We say that* $\mathcal{N}$ *term captures* A *iff* $\tau^{\mathcal{N}}_{A,\nu}$ *exists for all cardinals* ν *of* $\mathcal{N}$*.*

If $\mathcal{N}, \Gamma$ are as in Definition 2.1 and $A \in \Gamma$, then [4] shows that $\mathcal{N}$ term captures A. Later on, if the context is clear, we'll simply say capture instead of term capture or Suslin capture. We now define A-iterability.

Definition 2.3. *Let* $\mathcal{N}, \Gamma$ *be as above and* $A \in \Gamma$*. We say that* $\mathcal{N}$ *is* A*-iterable if* $\mathcal{N}$ *term captures* A *and*

1. *for any maximal tree* $\mathcal{T}$ *on* $\mathcal{N}$*, there is a cofinal branch* b *such that the branch embedding* $i^{\mathcal{T}}_b =_{def} i$ *moves the term relation for* A *correctly i.e., for any* κ *cardinal in* $\mathcal{N}$*,* $i(\tau^{\mathcal{N}}_{A,\kappa}) = \tau^{\mathcal{M}^{\mathcal{T}}_b}_{A,i(\kappa)}$*;*
2. *if* $\mathcal{T}$ *on* $\mathcal{N}$ *is short, then there is a branch* b *such that* $\mathcal{Q}(b, \mathcal{T})$[7] *exists and* $\mathcal{Q}(b, \mathcal{T}) \trianglelefteq Lp^{\Gamma}(\mathcal{M}(\mathcal{T}))$[8]*.*

The above definition obviously generalizes to define $\vec{A}$-iterability for any finite sequence $\vec{A}$. Also, the definition of A-iterability can be phrased in terms of the existence of a quasi-strategy (for the good player) in a certain iteration game.

Definition 2.4. *Let* $\mathcal{N}$ *be* k*-suitable with respect to* Σ^2_1 *and* $k < \omega$*. Let* $\vec{A} = \langle A_i \mid i \leq n \rangle$ *be a sequence of OD sets of reals and* $\nu = (\delta_0^{+\omega})^{\mathcal{N}}$*. Then*

---

[6] Γ is a good pointclass if it is closed under recursive preimages, closed under $\exists^{\mathbb{R}}$, is ω-parametrized, and has the scale property

[7] $\mathcal{Q}(b, \mathcal{T})$ is called the $\mathcal{Q}$-structure and is defined to be the least initial segment of $\mathcal{M}^{\mathcal{T}}_b$ that defines the failure of Woodinness of $\delta(\mathcal{T})$.

[8] This implicitly assumes that $\mathcal{Q}(b, \mathcal{T})$ has no extenders overlapping $\delta(\mathcal{T})$. We're only interested in trees $\mathcal{T}$ arising from comparisons between suitable mice and for such trees, $\mathcal{Q}$ structures have no extenders overlapping $\delta(\mathcal{T})$.

1. $\gamma^{\mathcal{N}}_{\vec{A}} = \sup(\{\xi | \xi$ *is definable over* $(\mathcal{N}|\nu, \tau^{\mathcal{N}}_{A_0,\delta_k}, ..., \tau^{\mathcal{N}}_{A_n,\delta_k})\}) \cap \delta_0$*;*
2. $H^{\mathcal{N}}_{\vec{A}} = Hull^{\mathcal{N}}(\gamma^{\mathcal{N}}_{\vec{A}} \cup \{\tau^{\mathcal{N}}_{A_0,\delta_k}, ..., \tau^{\mathcal{N}}_{A_n,\delta_k}\})$*, where we take the full elementary hull without collapsing.*

From now on, we will write $\tau^{\mathcal{N}}_{A}$ without further clarifying that this stands for $\tau^{\mathcal{N}}_{A,\delta}$ where δ is the largest Woodin cardinal of $\mathcal{N}$. Also, we'll occasionally say k-suitable without specifying the pointclass Γ.

Definition 2.5. *Let* $\mathcal{N}$ *be* k*-suitable with respectd to some pointclass* Γ *and* $A \in \Gamma$*.* $\mathcal{N}$ *is strongly* A*-iterable if* $\mathcal{N}$ *is* A*-iterable and for any suitable* $\mathcal{M}$ *such that if* $i, j : \mathcal{N} \to \mathcal{M}$ *are two* A*-iteration maps then* $i \restriction H^{\mathcal{N}}_{A} = j \restriction H^{\mathcal{M}}_{A}$*.*

Definition 2.6. *Let* Γ *be a good pointclass and* $\mathcal{N}$ *be* k*-suitable with respect to* Γ *for some* k*. Let* $\mathcal{A}$ *be a countable collection of sets of reals in* $\Gamma \cup \breve{\Gamma}$*. We say* $\mathcal{A}$ *guides a strategy* Σ *for* $\mathcal{N}$ *if whenever* $\mathcal{T}$ *is a countable, normal iteration tree on* $\mathcal{N}$ *by* Σ *of limit length, and* $b = \Sigma(\mathcal{T})$*, then*

1. *if* $\mathcal{T}$ *is short, then* $\mathcal{Q}(b, \mathcal{T})$ *exists and* $\mathcal{Q}(b, \mathcal{T}) \trianglelefteq Lp^{\Gamma}(\mathcal{M}(\mathcal{T}))$[9]*, and*
2. *if* $\mathcal{T}$ *is maximal, then* $i^{\mathcal{T}}_b(\tau^{\mathcal{N}}_{A,\mu}) = \tau^{\mathcal{M}^{\mathcal{T}}_b}_{A,i_b(\mu)}$ *for all* $A \in \mathcal{A}$ *and cardinals* $\mu \geq \delta^{\mathcal{N}}_k$ *of* $\mathcal{N}$ *and* $\delta = \sup\{\gamma^{\mathcal{M}^{\mathcal{T}}_b}_{A} \mid A \in \mathcal{A}\}$ *where* $\delta = i^{\mathcal{T}}_b(\delta_k)$*.*

The most important instance of the above definition used in this paper is when $\mathcal{A}$ is a self-justifying-system that seals a proper weak gap. A strategy guided by such an $\mathcal{A}$ has many desirable properties.

Finally, we quote two more important theorems of Woodin's that we'll be using repeatedly throughout the paper.

Theorem 2.7 (Woodin, Theorem 10.3 of [10])**.** *Assume* AD^+ *and suppose* Γ *is a good pointclass and is not the last good pointclass. There is then a function* F *defined on* $\mathbb{R}$ *such that for a Turing cone of* x*,* $F(x) = \langle \mathcal{N}^*_x, \mathcal{M}_x, \delta_x, \Sigma_x \rangle$ *such that*

1. $\mathcal{N}^*_x | \delta_x = \mathcal{M}_x | \delta_x$*,*
2. $\mathcal{N}^*_x \models$ *"*δ_x *is the only Woodin cardinal",*
3. Σ_x *is the unique iteration strategy of* $\mathcal{M}_x$*,*
4. $\mathcal{N}^*_x = L(\mathcal{M}_x, \Lambda)$ *where* Λ *is the restriction of* Σ_x *to stacks* $\vec{\mathcal{T}} \in \mathcal{M}_x$ *that have finite length and are based on* $\mathcal{M}_x \restriction \delta_x$*,*

---

[9] Again we disregard the case where $\mathcal{Q}$-structures have overlapping extenders.

5. $(\mathcal{N}_x^*, \Sigma_x)$ *Suslin captures* Γ,

6. *for any* $\alpha < \delta_x$ *and for any* $\mathcal{N}_x^*$*-generic* $g \subseteq Coll(\omega, \alpha)$, $(\mathcal{N}_x^*[g], \Sigma_x)$ *Suslin captures* $Code((\Sigma_x)_{\mathcal{M}_x \restriction \alpha})$ *and its complement at* δ_x^+.

Theorem 2.8 (Woodin, unpublished but see [12])**.** *Assume* $AD^+ + V = L(\mathcal{P}(\mathbb{R}))$*. Suppose* A *is a set of reals such that there is a Suslin cardinal in the interval* $(w(A), \theta_A)$*. Then*

1. *For any real* x *the pointclass* $\Sigma_1^2(A, x)$ *has the scale property and hence,* $\Sigma_1^2(A)$ *is a good pointclass.*

2. $\delta_1^2(A)$ *is a Suslin cardinal and is the largest one below* θ_A.

3. $M_{\underset{\sim}{\Delta}_1^2(A)} \prec_{\Sigma_1} L(\mathcal{P}(\mathbb{R}))$.

4. $L_\Theta(\mathcal{P}(\mathbb{R})) \prec_{\Sigma_1} L(\mathcal{P}(\mathbb{R}))$.

2.2 The computation

We start with quoting a theorem of Steel's which essentially says that Mouse Capturing implies Mouse Capturing for mice over $\mathbb{R}$.

Theorem 2.9 (Steel, see [10])**.** *Assume* $AD^+ + V = L(\mathcal{P}(\mathbb{R})) + \Theta = \theta_0 + MC$[10]*. Then every set of reals is in a (countably iterable) mouse over* $\mathbb{R}$ *projecting to* $\mathbb{R}$*. In other words,* $V = K(\mathbb{R})$*, where*

$$K(\mathbb{R}) = L(\cup\{M \mid M \textit{ is sound premouse over } \mathbb{R},\ \rho_\omega(M) = \mathbb{R}, \textit{ and } M \textit{ is countably iterable}\}).$$

Next, we prove the following theorem of Woodin's which roughly states that HOD is coded into a subset of Θ.

Theorem 2.10. *Assume* $AD^+ + V = L(\mathcal{P}(\mathbb{R}))$*. Then* $HOD = L[P]$ *for some* $P \subseteq \Theta$ *in* HOD*.*

Proof. First, let

$$\mathbb{P} = \{(\vec{\alpha}, \vec{a}) \mid \vec{\alpha} = \langle \alpha_0, \alpha_1, ..., \alpha_n \rangle \in \Theta^{<\omega}, \vec{a} = \langle a_0, a_1, ..., a_n \rangle, \forall i \leq n (a_i \subseteq \alpha_i)\}.$$

---

[10] MC stands for Mouse Capturing, which states that if $x, y \in \mathbb{R}$ and $x \in OD(y)$ then x is in a sound mouse over y projecting to y.

$\mathbb{P}$ is a poset with the (obvious) order by extension. If g is a $\mathbb{P}$-generic over V then g induces an enumeration of order type ω of $(\Theta, \cup_{\gamma<\Theta}\mathcal{P}(\gamma))$. Now let

$$\mathbb{Q}^* = \{(\vec{\alpha}, A) \mid \vec{\alpha} = \langle \alpha_0, \alpha_1, ..., \alpha_n\rangle \in \Theta^{<\omega}, A \subseteq \mathcal{P}(\alpha_0) \times \mathcal{P}(\alpha_1) \times ... \times \mathcal{P}(\alpha_n), A \in OD\}.$$

The ordering on $\mathbb{Q}^*$ is defined as follows:

$$(\vec{\alpha}, A) \leq (\vec{\beta}, B) \Leftrightarrow \forall i < \text{dom}(\vec{\alpha})\vec{\alpha}(i) = \vec{\beta}(i), B|(\mathcal{P}(\vec{\alpha}(0)) \times ... \times \mathcal{P}(\vec{\alpha}(\text{dom}(\alpha) - 1))) \subseteq A.$$

There is a poset $\mathbb{Q} \in HOD \cap \mathcal{P}(\Theta)$ that is isomorphic to $\mathbb{Q}^*$ via an OD map π. For our convenience, whenever $p \in \mathbb{Q}$, we will write p^* for $\pi(p)$. Furthermore, we can define π so that elements of $\mathbb{Q}$ have the form $(\vec{\alpha}, A)$ whenever $p^* = (\vec{\alpha}, A^*)$. In other words, we can think of π as a bijection of Θ and the set of OD subsets of $\mathcal{P}(\alpha_0) \times \mathcal{P}(\alpha_1) \times ... \times \mathcal{P}(\alpha_n)$ for $\alpha_0, \alpha_1, ..., \alpha_n < \Theta$. For notational simplicity, if $p^* = (\vec{\alpha}, A^*)$, we write $o(p^*)$ for $\vec{\alpha}$ and $s(p^*)$ for A^*.

Claim 1. *Let g be $\mathbb{P}$-generic over V. Then g induces a $\mathbb{Q}$-generic over HOD G_g.*

Proof. As mentioned above, g induces a generic enumeration f of $(\Theta, \cup_{\gamma<\Theta}\mathcal{P}(\gamma))$ of order type ω. Furthermore, for each $n < \omega$, $f(n)_0 < \Theta$ and $f(n)_1 \subseteq f(n)_0$. Let

$$G = \cup_{n<\omega}\{(\langle f(0)_0, ..., f(n)_0\rangle, A) \in \mathbb{Q} \mid \langle f(0)_1, ..., f(n)_1\rangle \in A^*\}.$$

We claim that G is $\mathbb{Q}$-generic over HOD. To see this, let $D \subseteq \mathbb{Q}$, $D \in HOD$ be a dense set. Let $p = f|(n+1)$ for some n. It's enough to find a $q = (\langle \alpha_0, ..., \alpha_m\rangle, \langle a_0, ..., a_m\rangle) \in \mathbb{P}$ extending p such that $D_q \cap D \neq \emptyset$ where

$$D_q = \{(\langle \alpha_0, ..., \alpha_m\rangle, A) \mid \langle a_0, ..., a_m\rangle \in A^*\}.$$

If no such q exists, let

$$r = (\langle f(0)_0, ..., f(n)_0\rangle, B),$$

where $b \in B^* \Leftrightarrow \forall t \in D(b \notin s(t))$. Then r is a condition in $\mathbb{Q}$ with no extension in D. Contradiction. □

Claim 2. *Let G be a $\mathbb{Q}$-generic over HOD. Then G induces a filter g_G for $\mathbb{P}$.*

Proof. It's enough to define the range of the function f induced by G. For each $\alpha < \Theta$, $n < \omega$, and $\langle \alpha_0, ..., \alpha_n\rangle$, let $A_{\alpha,\langle\alpha_0,...,\alpha_n\rangle} = (\langle \alpha_0, ..., \alpha_n\rangle, A)$ such that $\forall a \in A^*(\alpha \in a(n))$. Then for each $\alpha < \Theta$, $n < \omega$, let $f(n)_0 = o(p^*)(n)$ for some $p \in G$ (this is well-defined) and

$$\alpha \in f(n)_1 \Leftrightarrow A_{\alpha,\langle f(0)_0,\ldots,f(n)_0\rangle} \in G.$$

This function f clearly induces a filter g_G on $\mathbb{P}$. $\square$

The proof of Claim 2 suggests that we can define a canonical term in HOD for a generic enumeration of $\cup_{\gamma<\Theta}\mathcal{P}(\gamma)$. Indeed, let $A_{\alpha,\langle\alpha_0,\ldots,\alpha_n\rangle}$ be as in the proof of Claim 2. For each $n < \omega$, let $\sigma_n = \{(p, \check{\alpha}) \mid p \in \mathbb{Q}, p \leq A_{\alpha,\langle\alpha_0,\ldots,\alpha_n\rangle}$ for some $\langle\alpha_0, \ldots, \alpha_n\rangle \in \Theta^{n+1}\}$; let $\tau = \{(A, \sigma_n) \mid n < \omega, A \in \mathbb{Q}\}$. Then it's easy to see that whenever G is $\mathbb{P}$-generic over HOD induced by a $\mathbb{P}$-generic over V, τ_G enumerates $\cup_{\gamma<\Theta}\mathcal{P}(\gamma)$ in order type ω. This means we can recover $\mathcal{P}(\mathbb{R})^V$ in the model $L[\mathbb{Q}, \tau][G]$ by AD^+ (here we only use the fact that every set of reals has an ∞-Borel code which is a bounded subset of Θ).

To sum up, we have $L[\mathbb{Q}, \tau] \subseteq HOD \subseteq L[\mathbb{Q}, \tau][G]$ for some $\mathbb{Q}$-generic over HOD G. By a standard argument, this implies that $L[\mathbb{Q}, \tau] = HOD$. $\square$

Now let $\mathcal{F} = \{(\mathcal{M},\vec{A}) \mid \vec{A}$ is a finite sequence of OD sets of reals and $\mathcal{M}$ is k-suitable for some k and is strongly $\vec{A}$-iterable$\}$. We say $(\mathcal{M}, \vec{A}) \leq_{\mathcal{F}} (\mathcal{N}, \vec{B})$ if $\vec{A} \subseteq \vec{B}$ and $\mathcal{M}$ iterates to a suitable initial segment of $\mathcal{N}$. We then let $\pi_{(\mathcal{M},\vec{A}),(\mathcal{N},\vec{B})} : H^{\mathcal{M}}_{\vec{A}} \to H^{\mathcal{N}}_{\vec{B}}$ be the unique map. $\mathcal{F}$ is directed system because given any $(\mathcal{M}, \vec{A}), (\mathcal{N}, \vec{B}) \in \mathcal{F}$, we can compare them using their iteration strategies. $\mathcal{F} \neq \varnothing$ by the following theorem of Woodin, which we sketch a proof of.

Theorem 2.11. *Assume $V = L(\mathcal{P}(\mathbb{R})) + AD^+ + MC + \Theta = \theta_0$. Given any OD set of reals A and any $n \in \omega$, there is an n-suitable M that is strongly A-iterable.*

Proof. We'll prove the theorem for $n = 1$. The other cases are similar. So suppose not. By Theorem 2.9, $V = K(\mathbb{R})$. Then $V \vDash \phi$ where $\phi = (\exists\alpha)(\alpha$ is in a proper gap and $\mathrm{K}(\mathbb{R})|\alpha \vDash$ "$\mathrm{ZF}^- + \Theta$ exists $+ (\exists A)(A$ is OD and there is no 1-suitable strongly A-iterable mouse)".

Let $\delta < \underset{\sim}{\delta}^2_1$ be least such that $K(\mathbb{R})|\delta \vDash \phi$. Then it is easy to see that δ is a successor ordinal, say $\delta = \gamma + 1$. Furthermore, γ ends a proper weak gap, say $[\overline{\gamma}, \gamma]$ for some $\overline{\gamma} < \gamma$. Fix the least such A as above. Note that $\gamma < \underset{\sim}{\delta}^2_1$. By minimality of γ, we get a self-justifying-system $\langle A_i \mid i < \omega\rangle$ of $OD^{K(\mathbb{R})|\gamma}$ sets of reals in $K(\mathbb{R})|\gamma$ that seals the gap[11]. We may and do assume $A = A_0$. Let $\Gamma = \Sigma_1^{K(\mathbb{R})|\overline{\gamma}}$ and Ω a good scaled pointclass beyond $K(\mathbb{R})|(\gamma + 1)$, i.e. $\mathcal{P}(\mathbb{R})^{K(\mathbb{R})|(\gamma+1)} \subsetneq \underset{\sim}{\Delta}_{\Omega}$. Ω exists because $\gamma < \underset{\sim}{\delta}^2_1$. Let N^* be a coarse Ω-Woodin, fully iterable mouse. Such an N^* exists by [3] or by Theorem 2.7. In fact by Theorem 2.7, one can choose N^* that Suslin captures Ω and the sequence $\langle A_i \mid i < \omega\rangle$. Also by [3], there are club-in-ORD^{N^*} many Γ-Woodin cardinals in N^*. It can be shown that the $L[E]$ construction reaches a $\mathcal{P}$ such that $\mathcal{P}$ is 1-suitable with respect to Γ (hence has canonical terms for the A_i's) and $\mathcal{P} \vDash$ "δ_0 and δ_1 are Woodin cardinals" where δ_0 and δ_1 are the first two Γ-Woodin cardinals in N^*. Let Σ be the strategy for $\mathcal{P}$ induced by that of N^*. By lifting up to the background strategy and using term condensation for the self-justifying-system, we get that Σ is guided by $\langle A_i \mid i < \omega \rangle$, hence $(\mathcal{P}, \Sigma)$ is strongly A-iterable. But then $K(\mathbb{R})|\gamma \vDash$"$\mathcal{P}$ is strongly A-iterable." This is a contradiction. $\square$

---

[11] This means that for all i, $\neg A_i$ and a scale for A_i are in $\langle A_i \mid i < \omega\rangle$. Furthermore, the A_i's are cofinal in the Wadge hierarchy of $K(\mathbb{R})|\gamma$.

Lemma 2.12. *Let the direct limit of $\mathcal{F}$ under the maps $\pi_{(M,\vec{A}),(N,\vec{B})}$ be $\mathcal{M}_\infty$. Then $\mathcal{M}_\infty$ is wellfounded.*

Proof. We sketch a proof. Suppose not. Using Σ_1-reflection, there is an N a level of $K(\mathbb{R})$ below $\utilde{\delta}^2_1$ satisfying the statement "$ZF^- + DC + MC + \mathcal{M}_\infty$ is illfounded". We may assume N is the first such level. Let

$U = \{(x, \mathcal{M}) : \mathcal{M}$ is a sound x-mouse, $\rho_\omega(\mathcal{M}) = \{x\}$, and has an iteration strategy in $N\}$.

Since MC holds in N, U is a universal $(\Sigma^2_1)^N$-set. Let $A \in N$ be an OD set of reals witnessing ϕ. We assume that A has the minimal Wadge rank among the sets witnessing ϕ. Using the results of Chapter 3 of [1], we can get a $\vec{B} = \langle B_i : i < \omega \rangle$ which is a semiscale on U^c such that each $B_i \in N$.

Because MC holds and $\Gamma^* =_{def} \mathcal{P}(\mathbb{R})^N \subsetneq \utilde{\Delta}^2_1$, there is a real x such that there is a sound mouse $\mathcal{M}$ over x such that $\rho(\mathcal{M}) = x$ and $\mathcal{M}$ doesn't have an iteration strategy in N. Fix then such an $(x, \mathcal{M})$ and let Σ be the strategy of $\mathcal{M}$. Let Γ be a good pointclass such that $Code(\Sigma), \vec{B}, U, U^c \in \utilde{\Delta}_\Gamma$. Let F be as in Theorem 2.7 and let z be such that $(\mathcal{N}^*_z, \delta_z, \Sigma_z)$ Suslin captures $Code(\Sigma), \vec{B}, U, U^c$.

We let $\Phi = (\Sigma^2_1)^N$. We have that Φ is a good pointclass. Because $\vec{B}$ is Suslin captured by $\mathcal{N}^*_z$, we have $(\delta^+_z)^{\mathcal{N}^*_z}$-complementing trees $T, S \in \mathcal{N}^*_z$ which capture $\vec{B}$. Let κ be the least cardinal of $\mathcal{N}^*_z$ which, in $\mathcal{N}^*_z$ is $< \delta_z$-strong.

Claim 1. *$\mathcal{N}^*_z \vDash$ "κ is a limit of points η such that $Lp^{\Gamma^*}(\mathcal{N}^*_z|\eta) \vDash$ "η is Woodin".*

Proof. The proof is an easy reflection argument. Let $\lambda = \delta^+_z$ and let $\pi : M \to \mathcal{N}^*_z|\lambda$ be an elementary substructure such that

1. $T, S \in ran(\pi)$,
2. if $\text{cp}(\pi) = \eta$ then $V_\eta^{\mathcal{N}^*_z} \subseteq M$, $\pi(\eta) = \delta_z$ and $\eta > \kappa$.

By elementarity, we have that $M \vDash$ "η is Woodin". Letting $\pi^{-1}(\langle T, S\rangle) = \langle \bar{T}, \bar{S}\rangle$, we have that $(\bar{T}, \bar{S})$ Suslin captures the universal Φ set over M at $(\eta^+)^M$. This implies that M is Φ-full and in particular, $Lp^{\Gamma^*}(\mathcal{N}_z^*|\eta) \in M$. Therefore, $Lp^{\Gamma^*}(\mathcal{N}_z^*|\eta) \vDash$ "η is Woodin". The claim then follows by a standard argument. $\square$

Let now $\langle \eta_i : i < \omega\rangle$ be the first ω points $< \kappa$ such that for every $i < \omega$, $Lp^{\Gamma^*}(\mathcal{N}_z^*|\eta_i) \vDash$ "η_i is Woodin". Let now $\langle \mathcal{N}_i : i < \omega\rangle$ be a sequence constructed according to the following rules:

1. $\mathcal{N}_0 = L[\vec{E}]^{\mathcal{N}_z^*|\eta_0}$,
2. $\mathcal{N}_{i+1} = (L[\vec{E}][\mathcal{N}_i])^{\mathcal{N}_z^*|\eta_{i+1}}$.

Let $\mathcal{N}_\omega = \cup_{i<\omega}\mathcal{N}_i$.

Claim 2. *For every $i < \omega$, $\mathcal{N}_\omega \vDash$ "η_i is Woodin" and $\mathcal{N}_\omega|(\eta_i^+)^{\mathcal{N}_\omega} = Lp^{\Gamma^*}(\mathcal{N}_i)$.*

Proof. It is enough to show that

1. $\mathcal{N}_{i+1} \vDash$ "η_i is Woodin",
2. $\mathcal{N}_i = V_{\eta_i}^{\mathcal{N}_{i+1}}$,
3. $\mathcal{N}_{i+1}|(\eta_i^+)^{\mathcal{N}_{i+1}} = Lp^{\Gamma^*}(\mathcal{N}_i)$.

To show 1-3, it is enough to show that if $\mathcal{W} \trianglelefteq \mathcal{N}_{i+1}$ is such that $\rho_\omega(W) \leq \eta_i$ then the fragment of $\mathcal{W}$'s iteration strategy which acts on trees above η_i is in Γ^*. Fix then i and $\mathcal{W} \trianglelefteq \mathcal{N}_{i+1}$ is such that $\rho_\omega(W) \leq \eta_i$. Let ξ be such that the if $\mathcal{S}$ is the ξth model of the full background construction producing $\mathcal{N}_{i+1}$ then $\mathbb{C}(\mathcal{S}) = \mathcal{W}$. Let $\pi : \mathcal{W} \to \mathcal{S}$ be the core map. It is a fine-structural map but that it irrelevant and we surpass this point. The iteration strategy of $\mathcal{W}$ is the π-pullback of the iteration strategy of $\mathcal{S}$. Let then $\nu < \eta_{i+1}$ be such that $\mathcal{S}$ is the ξth model of the full background construction of $\mathcal{N}_x^*|\nu$. To determine the complexity of the induced strategy of $\mathcal{S}$ it is enough to determine the strategy of $\mathcal{N}_x^*|\nu$ which acts on non-dropping stacks that are completely above η_i. Now, notice that by the choice of η_{i+1}, for any non-dropping tree $\mathcal{T}$ on $\mathcal{N}_x^*|\nu$ which is above η_i and is of limit length, if $b = \Sigma(\mathcal{T})$ then $\mathcal{Q}(b, \mathcal{T})$ exists and $\mathcal{Q}(b, \mathcal{T})$ has no overlaps, and $\mathcal{Q}(b, \mathcal{T}) \trianglelefteq Lp^{\Gamma^*}(\mathcal{M}(\mathcal{T}))$. This observation indeed shows that the fragment of the iteration strategy of $\mathcal{N}_x^*|\nu$ that acts on non-dropping stack that are above η_i is in Γ^*. Hence, the strategy of $\mathcal{W}$ is in Γ^*. $\square$

We now claim that there is $\mathcal{W} \trianglelefteq Lp(\mathcal{N}_\omega)$ such that $\rho(W) < \eta_\omega$. To see this suppose not. It follows from MC that $Lp(\mathcal{N}_\omega)$ is Σ_1^2-full. We then have that x is generic over $Lp(\mathcal{N}_\omega)$ at the extender algebra of $\mathcal{N}_\omega$ at η_0. Because $Lp(\mathcal{N}_\omega)[x]$ is Σ_1^2-full, we have that $\mathcal{M} \in Lp(\mathcal{N}_\omega)[x]$

and $Lp(\mathcal{N}_\omega) \vDash$ "$\mathcal{M}$ is η_ω-iterable" by fullness of $Lp(\mathcal{N}_\omega)$. Let $\mathcal{S} = (L[\vec{E}][x])^{\mathcal{N}_\omega[x]|\eta_2}$ where the extenders used have critical point $> \eta_0$. Then working in $\mathcal{N}_\omega[x]$ we can compare $\mathcal{M}$ with $\mathcal{S}$. Using standard arguments, we get that $\mathcal{S}$ side doesn't move and by universality, $\mathcal{M}$ side has to come short (see [3]). This in fact means that $\mathcal{M} \trianglelefteq \mathcal{S}$. But the same argument used in the proof of Claim 2 shows that every $\mathcal{K} \trianglelefteq \mathcal{S}$ has an iteration strategy in Γ^*, contradiction!

Let now $\mathcal{W} \trianglelefteq Lp(\mathcal{N}_\omega)$ such that $\rho_\omega(\mathcal{W}) < \eta_\omega$. Let k, l be such that $\rho_l(\mathcal{W}) < \eta_k$. We can now consider $\mathcal{W}$ as a $\mathcal{W}|\eta_k$-mouse and considering it such a mouse we let $\mathcal{N} = \mathbb{C}_l(\mathcal{W})$. Thus, $\mathcal{N}$ is sound above η_k. We let $\langle \gamma_i : i < \omega \rangle$ be the Woodin cardinals of $\mathcal{N}$ and $\gamma = \sup_{i<\omega} \gamma_i$.

Let Λ be the strategy of $\mathcal{N}$. We claim that Λ is Γ^*-fullness preserving above γ_k. To see this fix $\mathcal{N}^*$ which is a Λ-iterate of $\mathcal{N}$ such that the iteration embedding $i : \mathcal{N} \to \mathcal{N}^*$ exists. If $\mathcal{N}^*$ isn't Γ^*-full then there is a strong cutpoint ν of $\mathcal{N}^*$ and a $\mathcal{N}^*|\nu$-mouse $\mathcal{W}$ with iteration strategy in Γ^* such that $\rho_\omega(\mathcal{W}) = \nu$ and $\mathcal{W} \ntrianglelefteq \mathcal{N}^*$. If $\mathcal{N}^*$ is not sound above ν then $\mathcal{N}^*$ wins the coiteration with $\mathcal{W}$; but this then implies $\mathcal{W} \ntriangleleft \mathcal{N}^*$, which contradicts our assumption. Otherwise, $\mathcal{N}^* \triangleleft \mathcal{W}$, which is also a contradiction. Hence Λ is Γ^*-fullness preserving.

We can then realize N as a derived model of a Λ-iterate of N which extends a Prikry generic over N (the details of this construction can be found in [11]). From this, the proof of the lemma is the same as that in [8]. □

$\mathcal{M}_\infty$ has ω Woodin cardinals $(\delta_0^{\mathcal{M}_\infty}, \delta_1^{\mathcal{M}_\infty}...)$ cofinal in its ordinals. Woodin has shown the following.

Lemma 2.13 (section 7.7.2 of [4]). *$\theta_0 = \delta_0^{\mathcal{M}_\infty}$ and $HOD|\theta_0 = \mathcal{M}_\infty|\delta_0^{\mathcal{M}_\infty}$.*

We'll attempt to extend this computation to the full HOD. Now we define a strategy Σ_∞ for $\mathcal{M}_\infty$. For each $A \in OD \cap \mathcal{P}(\mathbb{R})$, let $\tau_{A,k}^{\mathcal{M}_\infty}$ = common value of $\pi_{(\mathcal{P},A),\infty}(\tau_{A,k}^{\mathcal{P}})$ where $\pi_{(\mathcal{P},A),\infty}$ is the direct limit map and $\tau_{A,k}^{\mathcal{P}}$ is the standard term of $\mathcal{P}$ that captures A at δ_k^P. Σ_∞ will be defined (in V) for (finite stacks of) trees on $\mathcal{M}_\infty|\delta_0^{\mathcal{M}_\infty}$ in $\mathcal{M}_\infty$. For $k \geq n$, $\mathcal{M}_\infty \vDash$ "$Col(\omega, \delta_n^{\mathcal{M}_\infty}) \times Col(\omega, \delta_k^{\mathcal{M}_\infty}) \Vdash (\tau_{A,n}^{\mathcal{M}_\infty})_g = (\tau_{A,k}^{\mathcal{M}_\infty})_h \cap \mathcal{M}_\infty[g]$" where g is $Col(\omega, \delta_n^{\mathcal{M}_\infty})$ generic and h is $Col(\omega, \delta_k^{\mathcal{M}_\infty})$ generic. This is just saying that the terms cohere with one another.

Let G be $Col(\omega, \lambda^{\mathcal{M}_\infty})$ generic over $\mathcal{M}_\infty$ where $\lambda^{\mathcal{M}_\infty}$ is the sup of Woodin cardinals in $\mathcal{M}_\infty$. Then $\mathbb{R}_G^*$ is the symmetric reals and $A_G^* := \cup_k (\tau_{A,k}^{\mathcal{M}_\infty})_{G|\delta_k^{\mathcal{M}_\infty}}$.

Proposition 2.14. *For all $A \subseteq \mathbb{R}$, A is OD, $L(A_G^*, \mathbb{R}_G^*) \vDash AD^+$.*

Proof. If not, by Σ_1-reflction, there is an $N = K(\mathbb{R})|\gamma$, where $\gamma < \underset{\sim}{\delta}_1^2$ such that N satisfies the statement "$ZF^- + DC + MC + \exists A(A$ is OD and $L(A_G^*, \mathbb{R}_G^*) \nvDash AD^+))$". We assume N is the least such. By the same construction as in Lemma 2.12, there is the next mouse $(\mathcal{N}, \Lambda)$ with derived model N.

In N, let $A \subseteq \mathbb{R}$ be the least OD set such that $L(A^*_G, \mathbb{R}^*_G) \nvDash AD$. Then there is an iterate $\mathcal{M}$ of $\mathcal{N}$ having preimages of all the terms $\tau^{\mathcal{M}_\infty}_{A,k}$. We may assume $\mathcal{M}$ has derived model N. Since we have AD^+, $\mathcal{M}$ thinks that its derived model satisfies that $L(A, \mathbb{R}) \vDash AD^+$. Now iterate $\mathcal{M}$ to $\mathcal{P}$ such that $\mathcal{M}_\infty$ is a proper initial segment of $\mathcal{P}$. By elementarity $L(A^*_G, \mathbb{R}^*_G) \vDash AD^+$. This is a contradiction. □

Definition 2.15. *Given a normal tree $\mathcal{T} \in \mathcal{M}_\infty$ and $\mathcal{T}$ is based on $\mathcal{M}_\infty|\theta_0$. $\mathcal{T}$ is by Σ_∞ if the following hold (the definition is similar for finite stacks):*

- *If $\mathcal{T}$ is short then Σ picks the branch guided by $\mathcal{Q}$-structure (as computed in $\mathcal{M}_\infty$).*
- *If $\mathcal{T}$ is maximal then $\Sigma_\infty(\mathcal{T})$ = the unique cofinal branch b which moves $\tau^{\mathcal{M}_\infty}_{A,0}$ correctly for all $A \in OD \cap \mathcal{P}(\mathbb{R})$ i.e. for each such A, $i_b(\tau^{\mathcal{M}_\infty}_{A,0}) = \tau^{\mathcal{M}^{\mathcal{T}}_b}_{A^*,0}$.*

Lemma 2.16. *Given any such $\mathcal{T}$ as above, $\Sigma_\infty(\mathcal{T})$ exists.*

Proof. Suppose not. By reflection (Theorem 2.8), there is a (least) $\gamma < \underset{\sim}{\delta}^2_1$ such that $N =_{def} K(\mathbb{R})|(\gamma) \vDash \phi$ where ϕ is the statement "$ZF^- + DC + MC + \exists \mathcal{T}(\Sigma_\infty(\mathcal{T})$ doesn't exist)". We have a self-justifying-system $\vec{B}$ for $\Gamma^* = \mathcal{P}(\mathbb{R})^N$. By the construction of Lemma 2.12, there exists a mouse $\mathcal{N}$ with ω Woodin cardinals which has strategy Γ guided by $\vec{B}$. By results of [8], $\mathcal{M}^N_\infty$ is an iterate of $\mathcal{N}$ by Γ below its first Woodin. By reflecting to a countable hull, it's easy to see that Σ^N_∞ is a tail of Γ (the reflection is just to make all relevant objects countable). But this is a contradiction because then Σ^N_∞ can pick a branch for $\mathcal{T}$ that moves all the term relations correctly. □

It is evident that $L(\mathcal{M}_\infty, \Sigma_\infty) \subseteq HOD$. Next, we show $\mathcal{M}_\infty$ and Σ_∞ capture all of HOD. In $L(\mathcal{M}_\infty, \Sigma_\infty)$, first construct (using Σ_∞) a mouse $\mathcal{M}^+_\infty$ extending $\mathcal{M}_\infty$ such that $\mathrm{o}(\mathcal{M}_\infty)$ is the largest cardinal of $\mathcal{M}^+_\infty$ as follows:

1. Let $\mathbb{R}^*_G$ be the symmetric reals obtained from a generic G over $\mathcal{M}_\infty$ of $\mathrm{Col}(\omega, < \lambda^{\mathcal{M}_\infty})$.
2. For each A^*_G (defined as above where $A \in \mathcal{P}(\mathbb{R}) \cap OD^{K(\mathbb{R})}$) (we know $L(\mathbb{R}^*_G, A^*_G) \vDash AD^+$), pull back the R^*_G-mice in this model to mice $\mathcal{S}$ extending $\mathcal{M}_\infty$ with $D(\mathcal{S}, \lambda^{\mathcal{M}_\infty}) = L(\mathbb{R}^*_G, A^*_G)$. For more on this translation technique now called S-constructions, see [3] or [11].
3. Let $\mathcal{M}^+_\infty = \cup_{\mathcal{S}} \mathcal{S}$ for all such $\mathcal{S}$ as above. $\mathcal{M}^+_\infty$ is independent of G. By a reflection argument like that in Lemma 2.12 and using a Prikry-like forcing (see [11]), we get that the translated mice over $\mathcal{M}_\infty$ are all compatible, no levels of $\mathcal{M}^+_\infty$ projects across $o(\mathcal{M}_\infty)$, and $\mathcal{M}^+_\infty$ contains as its initial segments all restriction of $\mathbb{R}^*_G$-mice in

$D(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty})$ (in fact, in the reflected universe N in Lemma 2.12, $\mathcal{M}_\infty^+$ is an iterate of the next mouse $\mathcal{N}$). This is just to say that $\mathcal{M}_\infty^+$ contains enough mice to compute HOD.

Remark 2.17. $\delta_0^{\mathcal{M}_\infty}$ is not collapsed by Σ_∞ because it is a cardinal in HOD. Σ_∞ is used to obtain the A_G^* above by moving correctly the $\tau_{A,0}^{\mathcal{M}_\infty}$ in genericity iterations. $L(\mathcal{M}_\infty)$ generally does not see the sequence $\langle \tau_{A,k}^{\mathcal{M}_\infty} \mid k \in \omega\rangle$ hence can't construct A_G^*. Also since Σ_∞ collapses $\delta_1^{\mathcal{M}_\infty}, \delta_2^{\mathcal{M}_\infty}$... (via genericity iterations), it doesn't make sense to talk about $D(L(\mathcal{M}_\infty, \Sigma_\infty))$.

Lemma 2.18. *$HOD \subseteq L(\mathcal{M}_\infty, \Sigma_\infty)$*

Proof. Using Theorem 2.10, we know $HOD = L[P]$ for some $P \subseteq \Theta$. Therefore, it is enough to show P $\in L(\mathcal{M}_\infty, \Sigma_\infty)$. Let ϕ be a formula defining P, i.e.

$$\alpha \in P \Leftrightarrow K(\mathbb{R}) \vDash \phi[\alpha].$$

Here we suppress the ordinal parameter. Now in $L(\mathcal{M}_\infty, \Sigma_\infty)$ let $\pi : \mathcal{M}_\infty|((\delta_0^{\mathcal{M}_\infty})^{++})^{\mathcal{M}_\infty} \to (\mathcal{M}_\infty)^{D(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty})}$ where π is according to Σ_∞. We should note that Σ_∞-iterates are cofinal in the directed system $\mathcal{F}$ defined in $D(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty})$ by the method of generic comparisons (see [3] for more on this).

Claim: $K(\mathbb{R}) \vDash \phi[\alpha] \Leftrightarrow D(M_\infty^+, \lambda^{M_\infty}) \vDash \phi[\pi(\alpha)]$ $(**)$

Proof. Otherwise, reflect the failure of $(**)$ as before to the least $K(\mathbb{R})|\gamma$ and get a self-justifying-system $\vec{B}$ along with an ω-suitable mouse $\mathcal{N}$ with $\vec{B}$-guided iteration strategy Γ. By genericity iteration above its first Woodin, we may assume $D(\mathcal{N}, \lambda^{\mathcal{N}}) = K(\mathbb{R})|\gamma$. Fix an α witnessing the failure of $(**)$. Let $\sigma : \mathcal{N}|((\delta_0^{\mathcal{N}})^{++})^{\mathcal{N}} \to (\mathcal{M}_\infty)^{D(\mathcal{N}, \lambda^{\mathcal{N}})}$ be the direct limit map by Γ (we may assume everything is countable to make sense of iterability). We may assume there is an $\overline{\alpha}$ such that $\sigma(\overline{\alpha}) = \alpha$. Notice here that $\Sigma_\infty^{K(\mathbb{R})|\gamma}$ is a tail of Γ as $\Sigma_\infty^{K(\mathbb{R})|\gamma}$ moves all the term relations for $OD^{K(\mathbb{R})|\gamma}$ sets of reals correctly and Γ is guided by the self-justifying system $\vec{B}$, which is cofinal in $\mathcal{P}(\mathbb{R}) \cap OD^{K(\mathbb{R})|\gamma}$. It then remains to see that:

$$D(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty}) \vDash \phi[\pi(\alpha)] \Leftrightarrow D(\mathcal{N}, \lambda^{\mathcal{N}}) \vDash \phi[\sigma(\overline{\alpha})] \quad (***).$$

To see that $(***)$ holds, we need to see that the fragment of Γ that defines $\sigma(\overline{\alpha})$ can be defined in $D(\mathcal{N}, \lambda^{\mathcal{N}})$. This then will give the equivalence in $(***)$. Because $\alpha < \delta_0^{\mathcal{M}_\infty^{K(\mathbb{R})|\gamma}} = \delta_0^{\mathcal{M}_\infty^{D(\mathcal{N}, \lambda^{\mathcal{N}})}}$, pick an $A \in \mathcal{P}(\mathbb{R}) \cap OD^{D(\mathcal{N}, \lambda^{\mathcal{N}})}$ such that $\gamma_{A,0}^{D(\mathcal{N}, \lambda^{\mathcal{N}})} > \alpha$. Then the fragment of Γ that defines $\sigma(\overline{\alpha})$ is definable from A (and $\mathcal{N}|(\delta_0^N)$) in $D(\mathcal{N}, \lambda^{\mathcal{N}})$, which is what we want.

The equivalence $(***)$ gives us a contradiction. □

The claim finishes the proof of $P \in L(\mathcal{M}_\infty, \Sigma_\infty)$ because the right hand side of the equivalence $(**)$ can be computed in $L(\mathcal{M}_\infty, \Sigma_\infty)$. This then implies $HOD = L[P] \subseteq L(\mathcal{M}_\infty, \Sigma_\infty)$. $\square$

Lemma 2.18 implies $HOD = L(\mathcal{M}_\infty, \Sigma_\infty)$, hence completes our computation. Let P be as in Theorem 2.10 and let ϕ be the Σ_1 formula defining P.

Remark 2.19. Woodin (unpublished) has also computed the full HOD for models satisfying $V = L(\mathcal{P}(\mathbb{R})) + AD^+ + \Theta = \theta_0$. To the best of the author's knowledge, here's a very rough idea of his computation. Let $\mathcal{M}_\infty, \Sigma_\infty, P$ be as above. For each $\alpha < \Theta$, let Σ_α be the fragment of Σ_∞ that moves α along the good branch of a maximal tree. Woodin shows that the structure $(\mathbb{R}^*_G, \langle \Sigma_\alpha \mid \alpha < \Theta \rangle)$ can compute the set P. This then gives us that $HOD \subseteq L(\mathcal{M}_\infty, \Sigma_\infty)$.

3 The $\Theta = \theta_{\alpha+1}$ case

3.1 A theory of hod mice

In this subsection, we summarize definitions and facts about hod mice that will be used in our computation in the next two sections. For basic definitions and notations that we omit, see [3].

We say a transitive set a is self-wellordered if there is a well-ordering of a in $L_1[a]$.

Definition 3.1. *Suppose Σ is an iteration strategy with hull-condensation, a is a transitive self-wellordered set such that $\mathcal{M}_\Sigma \in a$, $\kappa = (|a|^+)$, and Σ is a κ-iteration strategy. Then $Lp^\Sigma_\kappa(a) = \cup_{\alpha<\kappa} Lp^\Sigma_\alpha(a)$ where*

1. *$Lp^\Sigma_0(a) = a \cup \{a\}$*
2. *$Lp^\Sigma_{\alpha+1}(a) = \cup\{\mathcal{N} : \mathcal{N}$ is a sound $\kappa+1$-iterable Σ-mouse over $Lp^\Sigma_\alpha(a)$ projecting to $o(Lp^\Sigma_\alpha(a))\}$.*
3. *$Lp^\Sigma_\lambda(a) = \cup_{\alpha<\lambda} Lp^\Sigma_\alpha(a)$ for limit λ.*

We let $Lp^\Sigma(a) = Lp^\Sigma_1(a)$.

Definition 3.2. *Suppose $\mathcal{P} = J_\alpha^{\vec{E},S,T,\langle \nu_\xi, \mu_\xi : \xi < \varsigma \rangle}(\mathcal{N})$ is a layered hybrid premouse*[12]*. $\mathcal{P}$ is a hod premouse below* $\mathrm{AD}_\mathbb{R}$ + *"θ is measurable" if (see Figure 1)*

---

[12] For more on hybrid structures, see [3]


$\delta_\lambda^{+\omega}$
$\Sigma_{<\lambda}$ mouse
$\delta_\lambda, \Sigma_{<\lambda}$
Σ_α mouse
$\delta_\alpha, \Sigma_\alpha$
δ_1, Σ_1
δ_0, Σ_0

Figure 1: Hod premouse.

1. $\mathcal{P} \vDash$ ZFC-*Replacement* +"$\langle \nu_\xi : \xi < \varsigma \rangle$ *is the sequence of Woodin cardinals and their limits enumerated in the increasing order"+"for every* $\xi < \varsigma$, ν_ξ *is a strong cutpoint",*
2. $\varsigma = \lambda + 1$ *for some* λ, *for every* $n < \omega$, $\mathcal{P} \vDash$ "ν_λ^{+n} *exists" and* $o(\mathcal{P}) = \sup_{n<\omega}(\nu_\lambda^{+n})^{\mathcal{P}}$,
3. $\mathcal{P} \vDash$ " *for every* $\xi < \varsigma$, T_ξ *is an* $(< o(\mathcal{P}), < o(\mathcal{P}))$*-iteration strategy for* $\mathcal{P}|\mu_\xi$ *below* ν_ξ *with hull-condensation",*
4. $\mathcal{P} \vDash \mathcal{P}|\mu_\xi \cong Lp_\omega^{S\oplus T_{<\xi}}(\mathcal{P} \restriction \nu_\xi)$ *and for limit* ξ, $\mathcal{P} \vDash \nu_\xi^+ = o(Lp^{S\oplus T_{<\xi}}(\mathcal{P} \restriction \nu_\xi))$,
5. $\mathcal{P} \vDash$ "*if* η *and* $\xi < \varsigma$ *are such that* $\eta \in (\nu_\xi, \nu_{\xi+1}]$ *then* $Lp^{S\oplus T_\xi}(\mathcal{P}|\eta) \trianglelefteq \mathcal{P}$",
6. *if for some* $\xi < \varsigma$, $\xi + 2 \leq \varsigma$ *then* $\mathcal{P} \vDash$ " *for any cardinal* $\eta \in (\nu_\xi, \nu_{\xi+1})$, $\mathcal{P}|\eta$ *is* $(< o(\mathcal{P}), < o(\mathcal{P}))$*-iterable for non-dropping trees via a strategy consistent with* T".

It is the requirement that ν_ξ is a strong cutpoint that makes our hod premice "below $\mathrm{AD}_{\mathbb{R}}$ + "θ is measurable"". From now on we drop "below $\mathrm{AD}_{\mathbb{R}}$ + "θ is measurable"" and simply say "$\mathcal{P}$ is a hod premouse". Also, we let "$\mathcal{P}$ is a hod premouse" stand for hod premmice over $\emptyset$, i.e., $S = \emptyset$ and $\mathcal{N} = \emptyset$. Clearly all notions that we will define for lightface hod premice easily generalize to non-lightface hod premice. If $S \neq \emptyset$ and there is a strategy Λ such that $S = \mathcal{P} \cap \Lambda$ then we say $\mathcal{P}$ is a Λ-hod premouse. All the notions that we isolate for hod premice easily generalize to Λ-hod premice. If $\mathcal{P}$ is a hod premouse then we change our notation and let $\langle \delta_\xi, \mu_\xi : \xi \leq \lambda \rangle = \langle \nu_\xi, \mu_\xi : \xi \leq \varsigma - 1 \rangle$ and we let $\lambda^{\mathcal{P}} = \lambda$. We also let $\delta^{\mathcal{P}} = \delta^{\mathcal{P}}_{\lambda^{\mathcal{P}}}$. If in Definition 3.2, $\lambda^{\mathcal{P}}$ is a successor ordinal then we let $\mathcal{P}^- = \mathcal{P}|\mu_{\lambda^{\mathcal{P}}-1}$. If $\lambda^{\mathcal{P}}$ is limit then we let $\mathcal{P}^- = \mathcal{P}|\delta^{\mathcal{P}}$. For $\xi \leq \lambda$ we let $\mathcal{P}(\xi) = \mathcal{P}|\mu_\xi$. Also, for $\alpha < \lambda^{\mathcal{P}}$, we let Σ^P_α be the iteration strategy of $\mathcal{P}(\alpha)$ and for limit $\alpha \leq \lambda^{\mathcal{P}}$ we let $\Sigma^{\mathcal{P}}_{<\alpha} = \oplus_{\beta<\alpha}\Sigma_\beta$. We let $\Sigma^{\mathcal{P}} = \Sigma^{\mathcal{P}}_{<\lambda^{\mathcal{P}}}$ if $\lambda^{\mathcal{P}}$ is limit and $\Sigma^{\mathcal{P}} = \Sigma^{\mathcal{P}}_{\lambda^{\mathcal{P}}-1}$ if $\lambda^{\mathcal{P}}$ is a successor.

Definition 3.3. *$(\mathcal{P}, \Sigma)$ is a hod pair if $\mathcal{P}$ is a hod premouse and Σ is an iteration strategy for $\mathcal{P}$ with hull condensation.*

Theorem 3.4. *Suppose $(\mathcal{P}, \Sigma)$ is a hod pair such that Σ has branch condensation. Then Σ is pullback consistent, positional and commuting.*

The proof of Theorem 3.4 can be found in [3]. For hod pairs $(\mathcal{M}_\Sigma, \Sigma)$, if Σ is a strategy with branch condensation and $\vec{\mathcal{T}}$ is a stack on M_Σ with last model N, $\Sigma_{N,\vec{\mathcal{T}}}$ is independent of $\vec{\mathcal{T}}$. Therefore, later on we will omit the subscript $\vec{\mathcal{T}}$ from $\Sigma_{N,\vec{\mathcal{T}}}$ whenever Σ is a strategy with branch condensation and $\mathcal{M}_\Sigma$ is a hod mouse.

Definition 3.5. *Suppose $\mathcal{P}$ and $\mathcal{Q}$ are two hod premice. Then $\mathcal{P} \trianglelefteq_{hod} \mathcal{Q}$ if there is $\alpha \leq \lambda^{\mathcal{Q}}$ such that $\mathcal{P} = \mathcal{Q}(\alpha)$.*

If $\mathcal{P}$ and $\mathcal{Q}$ are hod premice such that $\mathcal{P} \trianglelefteq_{hod} \mathcal{Q}$ then we say $\mathcal{P}$ is a hod initial segment of $\mathcal{Q}$. If $(\mathcal{P}, \Sigma)$ is a hod pair, $\mathcal{Q}$ is a non-dropping Σ-iterate of $\mathcal{P}$ and $\mathcal{R} \trianglelefteq_{hod} \mathcal{Q}$ then we let $\Sigma_{\mathcal{R}}$ be the iteration strategy for $\mathcal{R}$ given by $\Sigma_{\mathcal{Q}}$.

All hod pairs $(\mathcal{P}, \Sigma)$ have the property that Σ has hull condensation and therefore, mice relative to Σ make sense. To state the Strong Mouse Capturing we need to introduce the notion of Γ-fullness.

Definition 3.6. *Suppose Σ is an iteration strategy with hull-condensation, a is a transitive set such that $\mathcal{M}_\Sigma \in a$*[13] *and Γ is a pointclass closed under boolean operations and continuous images and preimages. Then $Lp^{\Gamma,\Sigma}_{\omega_1}(a) = \cup_{\alpha<\kappa} Lp^{\Gamma,\Sigma}_\alpha(a)$ where*

1. *$Lp^{\Gamma,\Sigma}_0(a) = a \cup \{a\}$*
2. *$Lp^{\Gamma,\Sigma}_{\alpha+1}(a) = \cup\{\mathcal{M} : \mathcal{M}$ is a sound $\Sigma_{\mathcal{Q}(\alpha),\mathcal{T}}$-mouse over $Lp^{\Gamma,\Sigma}_\alpha(a)$ projecting to η and having an iteration strategy in $\Gamma\}$.*
3. *$Lp^{\Gamma,\Sigma}_\lambda(a) = \cup_{\alpha<\lambda} Lp^{\Gamma,\Sigma}_\alpha(a)$ for limit λ.*

We let $Lp^{\Gamma,\Sigma}(a) = Lp^{\Gamma,\Sigma}_1(a)$

---

[13] $\mathcal{M}_\Sigma$ is the structure that Σ-iterates.

Definition 3.7 (Γ-Fullness preservation)**.** *Suppose* $(\mathcal{P}, \Sigma)$ *is a hod pair and* Γ *is a pointclass closed under boolean operations and continuous images and preimages. Then* Σ *is a* Γ*-fullness preserving if whenever* $(\vec{\mathcal{T}}, \mathcal{Q}) \in I(\mathcal{P}, \Sigma)$, $\alpha + 1 \leq \lambda^{\mathcal{Q}}$ *and* $\eta > \delta_\alpha$ *is a strong cutpoint of* $\mathcal{Q}(\alpha + 1)$, *then*

$$\mathcal{Q}|(\eta^+)^{\mathcal{Q}(\alpha+1)} = Lp^{\Gamma, \Sigma_{\mathcal{Q}(\alpha), \vec{\mathcal{T}}}}(\mathcal{Q}|\eta).$$

and

$$\mathcal{Q}|(\delta_\alpha^+)^{\mathcal{Q}} = Lp^{\Gamma, \oplus_{\beta<\alpha}\Sigma_{\mathcal{Q}(\beta+1), \vec{\mathcal{T}}}}(\mathcal{Q}(\alpha)).$$

Here is the statement of the strong mouse capturing.

Definition 3.8 (The Strong Mouse Capturing)**.** *The Strong Mouse Capturing (SMC) is the statement: Suppose* $(\mathcal{P}, \Sigma)$ *is a hod pair such that* Σ *has branch condensation and is* Γ*-fullness preserving for some* Γ*. Then for any* $x, y \in \mathbb{R}$, $x \in OD_\Sigma(y)$ *iff* x *is in some* Σ*-mouse over* $\langle \mathcal{P}, y \rangle$*.*

When $(\mathcal{P}, \Sigma) = \emptyset$ in the statement of Definition 3.8 we get the ordinary Mouse Capturing (MC). The Strong Mouse Set Conjecture (SMSC) just conjectures that SMC hods.

Definition 3.9 (Strong Mouse Set Conjecture)**.** *Assume* AD^+ *and that there is no mouse with a superstrong cardinal. Then* SMC *holds.*

Finally, we introduce the Γ-hod pair construction from [3] which is crucial for our HOD analysis. Suppose Γ is a pointclass closed under complements and under continuous preimages. Let

$$HP^\Gamma = \{(\mathcal{P}, \Lambda) \mid (\mathcal{P}, \Lambda) \text{ is a hod pair and } Code(\Lambda) \in \Gamma\},$$

and

$$\begin{aligned} Mice^\Gamma = \{(a, \Lambda, \mathcal{M}) \quad & \mid a \in HC,\ a \text{ is self-wellordered transitive, } \Lambda \text{ is an iteration} \\ & \text{strategy such that } (\mathcal{M}_\Lambda, \Lambda) \in HP^\Gamma,\ \mathcal{M}_\Lambda \in a, \text{ and } \mathcal{M} \trianglelefteq Lp^{\Gamma,\Lambda}(a)\}. \end{aligned}$$

If $\Gamma = \mathcal{P}(\mathbb{R})$, we let $HP = HP^\Gamma$ and $Mice = Mice^\Gamma$. Suppose $(\mathcal{M}_\Sigma, \Sigma) \in HP^\Sigma$. Let

$$Mice^\Gamma_\Sigma = \{(a, \mathcal{M}) \mid (a, \Sigma, \mathcal{M}) \in Mice^\Gamma\}.$$

Let $A \subseteq \mathbb{R}$ be such that $w(A) \geq w(\Gamma)$. Let A_Γ be the set of reals $\langle \sigma_0, \sigma_1 \rangle$ of continuous functions such that $\sigma_0^{-1}[A]$ is a code for a set in HP^Σ and $\sigma_1^{-1}[A]$ is a code for $(a, \Lambda, \mathcal{M}, \Psi)$ where $(a, \Lambda, \mathcal{M}) \in Mice^\Gamma$ and Ψ is the unique strategy of $\mathcal{M}$.

Definition 3.10 (Γ-hod pair construction)**.** *Let Γ, A be as above. Suppose (M, δ, Σ) is such that $M \models ZFC$ - Replacement, (M, δ) is countable, δ is an uncountable cardinal in M, Σ is an (ω_1, ω_1)-iteration strategy for M, $\Sigma \cap (L_1(V_\delta))^M \in M$. Suppose M locally Suslin, co-Suslin captures A_Γ. Then the Γ-hod pair construction of M below δ is a sequence $\langle\langle \mathcal{N}_\xi^\beta \mid \xi < \delta\rangle, \mathcal{P}_\beta, \Sigma_\beta, \delta_\beta \mid \beta \leq \Omega\rangle$ that satisfies the following properties.*

1. *$M \models$ "for all $\beta < \Omega, (\mathcal{P}_\beta, \Sigma_\beta)$ is a hod pair such that $\Sigma_\beta \in \Gamma$"*

2. *$\langle \mathcal{N}_\xi^0 \mid \xi < \delta\rangle$ are the models of the $L[\vec{E}]$-construction of V_δ^M and $\langle \mathcal{N}_\xi^\beta \mid \xi < \delta\rangle$ are the models of the $L[\vec{E}, \Sigma_\beta]$-construction of V_δ^M. δ_0 is the least γ such that $o(\mathcal{N}_\gamma^0) = \gamma$ and $Lp^\Gamma(\mathcal{N}_\gamma^0) \models$ "γ is Woodin" and $\delta_{\beta+1}$ is the least γ such that $o(\mathcal{N}_\gamma^{\beta+1}) = \gamma$ and $Lp^{\Gamma,\Sigma_\beta}(\mathcal{N}_\gamma^{\beta+1}) \models$ "γ is Woodin".*

3. *$\mathcal{P}_0 = Lp_\omega^\Gamma(\mathcal{N}_{\delta_0}^0)$ and Σ_0 is the canonical strategy of $\mathcal{P}_0$ induced by Σ.*

4. *Suppose $\delta_{\beta+1}$ exists, $\mathcal{N}_{\delta_{\beta+1}}^{\beta+1}$ doesn't project across δ_β. Furthermore, if $\beta = 0$ or is successor and $\mathcal{N}_{\delta_{\beta+1}}^{\beta+1} \models$ "δ_β is Woodin" and if β is limit then $(\delta_\beta^+)^{\mathcal{P}_\beta} = (\delta_\beta^+)^{\mathcal{N}_{\delta_{\beta+1}}^{\beta+1}}$, then $\mathcal{P}_{\beta+1} = Lp_\omega^{\Gamma,\Sigma_\beta}(\mathcal{N}_{\delta_{\beta+1}}^{\beta+1})$ and $\Sigma_{\beta+1}$ is the canonical strategy $\mathcal{P}_{\beta+1}$ induced by Σ.*

5. *For limit ordinals β, letting $\mathcal{P}_\beta^* = \cup_{\gamma<\beta}\mathcal{P}_\gamma$, $\Sigma_\beta^* = \oplus_{\gamma<\beta}\Sigma_\gamma$, and $\delta_\beta = sup_{\gamma<\beta}\delta_\gamma$, if δ_β is not measurable in M and $\delta_\beta < \delta$ then $\langle \mathcal{N}_\xi^\beta \mid \xi < \delta\rangle$ are the models of the $L[\vec{E}, \Sigma_\beta^*]$-construction of V_δ^M. If there isn't any γ such that $o(\mathcal{N}_\gamma^\beta) = \gamma$ and $Lp^{\Gamma,\Sigma_\beta^*}(\mathcal{N}_\gamma^\beta) \models$ "γ is Woodin" then we let $\mathcal{P}_\beta$ be undefined. Otherwise, let γ be the least such that $o(\mathcal{N}_\gamma^\beta) = \gamma$ and $Lp^{\Gamma,\Sigma_\beta^*}(\mathcal{N}_\gamma^\beta) \models$ "γ is Woodin." If $\mathcal{N}_\gamma^\beta$ doesn't project across δ_β then $\mathcal{P}_\beta = \mathcal{N}_\gamma^\beta|(\delta_\beta^{+\omega})^{\mathcal{N}_\gamma^\beta}$, and Σ_β is the canonical iteration strategy for $\mathcal{P}_\beta$ induced by Σ. Otherwise, let $\mathcal{P}_\beta$ be undefined.*

3.2 The computation

Recall that in this section we assume that $\Theta = \theta_{\alpha+1}$. We state without proof the theorem that will be important for the computation in this section.

Theorem 3.11. *Assume $AD^+ + SMC$. Suppose $(\mathcal{P}, \Sigma)$ is a hod pair below $AD_{\mathbb{R}}$ + "Θ is regular"[14] such that Σ has branch condensation and is fullness preserving. Then*

$$\{A \subseteq \mathbb{R} : A \in OD_\Sigma(y) \text{ for some real } y\} = \mathcal{P}(\mathbb{R}) \cap K^\Sigma(\mathbb{R}),$$

where

---

[14] i.e., there are no inaccessible limit of Woodins in $\mathcal{P}$.

$$K^{\Sigma}(\mathbb{R}) = \cup\{\mathcal{M} : \mathcal{M} \textit{ is a sound countably iterable } \Sigma\textit{-mouse over } \mathbb{R} \textit{ with } \rho_\omega(\mathcal{M}) = \mathbb{R}\}$$

.

Remark: Theorem 3.11 is a generalization of Theorem 2.9. The part that we do not spell out here is the precise definition of a Σ-mouse over $\mathbb{R}$. It turns out that it is quite a delicate issue (see [3] for more details) and is not important for us here.

First we need to compute V_Θ^{HOD}. Here's what is done in [3] regarding this computation.

Theorem 3.12 (Sargsyan, see Section 4.3 in [3]). *Let $\Gamma = \{A \subseteq \mathbb{R} \mid w(A) < \theta_\alpha\}$. Then there is a hod pair $(\mathcal{P}, \Sigma)$ such that*

1. *Σ is fullness preserving and has branch condensation;*
2. *$\Gamma(\mathcal{P}, \Sigma) = \Gamma$ where $\Gamma(\mathcal{P}, \Sigma) = \{A \subseteq \mathbb{R} \mid A \leq_w \Sigma_{\mathcal{Q}(\beta)}$ for some $\beta < \lambda^{\mathcal{Q}}$ where $\mathcal{Q}$is a $\Sigma-$ iterate of $\mathcal{P}\}$;*
3. *$\mathcal{M}^+_\infty(\mathcal{P}, \Sigma)|\theta_\alpha = V_\alpha^{HOD}$, where $\mathcal{M}^+_\infty(\mathcal{P}, \Sigma)$ is the direct limit of all Σ-iterates of $\mathcal{P}$.*

It is clear that there is no hod pair $(\mathcal{P}, \Sigma)$ satisfying Theorem 3.12 with Γ replaced by $\mathcal{P}(\mathbb{R})$ as this would imply that $\Sigma \notin V$. So to compute V_Θ^{HOD}, we need to mimic the computation in Section 2. First, we need to generalize the notion of suitability.

Definition 3.13 (n-suitable pair). *$(\mathcal{P}, \Sigma)$ is an n-suitable pair if there is δ such that $(\mathcal{P}|(\delta^{+\omega})^{\mathcal{P}}, \Sigma)$ is a hod pair and*

1. *$\mathcal{P} \models$ ZFC - Replacement + "there are n Woodin cardinals, $\eta_0 < \eta_1 < ... < \eta_{n-1}$ above δ";*
2. *$o(\mathcal{P}) = sup_{i<\omega}(\eta_{n-1})^{+i^P}$;*
3. *$\mathcal{P}$ is a Σ-mouse over $\mathcal{P}|\delta$;*
4. *for any $\mathcal{P}$-cardinal $\eta > \delta$, if η is a strong cutpoint then $\mathcal{P}|(\eta^+)^{\mathcal{P}} = Lp^{\Sigma}(\mathcal{P}|\eta)$.*

For $\mathcal{P}, \delta$ as in the above definition, let $\mathcal{P}^- = \mathcal{P}|(\delta^{+\omega})^{\mathcal{P}}$ and $\mathbb{B}(\mathcal{P}^-, \Sigma) = \{B \subseteq \mathcal{P}(\mathbb{R}) \times \mathbb{R} \times \mathbb{R} \mid B$ is OD, and for any $(\mathcal{Q}, \Lambda)$ iterate of $(\mathcal{P}^-, \Sigma)$, and for any $(x, y) \in B_{(\mathcal{Q},\Lambda)}, x$ codes $\mathcal{Q}\}$. Suppose $B \in \mathbb{B}(P^-, \Sigma)$ and $\kappa < o(\mathcal{P})$. Let $\tau^{\mathcal{P}}_{B,\kappa}$ be the canonical term in $\mathcal{P}$ that captures B at κ i.e. for any $g \subseteq Col(\omega, \kappa)$ generic over $\mathcal{P}$

$$B_{(\mathcal{P}^-,\Sigma)} \cap \mathcal{P}[g] = (\tau^{\mathcal{P}}_{B,\kappa})_g.$$

For each $m < \omega$, let

$$\gamma^{\mathcal{P},\Sigma}_{B,m} = \sup(Hull^{\mathcal{P}}(\tau^{\mathcal{P}}_{B,(\eta^{+m}_{n-1})^{\mathcal{P}}}) \cap \eta_0),$$

$$H^{\mathcal{P},\Sigma}_{B,m} = Hull^{\mathcal{P}}(\gamma^{\mathcal{P},\Sigma}_{B,m} \cup \{\tau^{\mathcal{P}}_{B,(\eta^{+m}_{n-1})^{\mathcal{P}}}\}),$$

$$\gamma^{\mathcal{P},\Sigma}_{B} = \sup_{m<\omega}\gamma^{\mathcal{P},\Sigma}_{B,m},$$

and

$$H^{\mathcal{P},\Sigma}_{B} = \cup_{m<\omega} H^{\mathcal{P},\Sigma}_{B,m}.$$

Similar definitions can be given for $\gamma^{\mathcal{P},\Sigma}_{\vec{B},m}, H^{\mathcal{P},\Sigma}_{\vec{B},m}, \gamma^{\mathcal{P},\Sigma}_{\vec{B}}, H^{\mathcal{P},\Sigma}_{\vec{B}}$ for any finite sequence $\vec{B} \in \mathbb{B}(\mathcal{P}^-, \Sigma)$. One just needs to include relevant terms for each element of $\vec{B}$ in each relevant hull. Now we define the notion of B-iterability.

Definition 3.14 (B-iterability)**.** *Let $(\mathcal{P}, \Sigma)$ be an n-suitable pair and $B \in \mathbb{B}(\mathcal{P}^-, \Sigma)$. We say $(\mathcal{P}, \Sigma)$ is B-iterable if for all $k < \omega$, player II has a winning quasi-strategy for the game $G^{(P,\Sigma)}_{B,k}$ defined as follows. The game consists of k rounds. Each round consists of a main round and a subround. Let $(\mathcal{P}_0, \Sigma_0) = (\mathcal{P}, \Sigma)$. In the main round of the first round, player I plays countable stacks of normal nondropping trees based on $\mathcal{P}^-_0$ or its images and player II plays according to Σ_0 or its tails. If the branches chosen by player II does not move some term for B correctly, he loses. Player I has to exit the round at a countable stage; otherwise, he loses. Suppose $(\mathcal{P}^*, \Sigma^*)$ is the last model after the main round is finished. In the subround, player I plays a normal tree above $(\mathcal{P}^*)^-$ or its images based on a window of two consecutive Woodins. Player II plays a branch that moves all terms for B correctly. Otherwise, he loses. Suppose $(\mathcal{P}_1, \Sigma_1)$ is the last model of the subround. If II hasn't lost, the next round proceeds the same way as the previous one but for the pair $(\mathcal{P}_1, \Sigma_1)$. If the game lasts for k rounds, II wins.*

Definition 3.15 (Strong B-iterability)**.** *Let $(\mathcal{P}, \Sigma)$ be an n-suitable pair and $B \in \mathbb{B}(\mathcal{P}^-, \Sigma)$. We say $(\mathcal{P}, \Sigma)$ is strongly B-iterable if $(\mathcal{P}, \Sigma)$ is B-iterable and if r_1 is a run of $G^{\mathcal{P},\Sigma}_{B,n_1}$ and r_1 is a run of $G^{\mathcal{P},\Sigma}_{B,n_2}$ for some $n_1, n_2 < \omega$ according to the winning quasi-strategy of $\mathcal{P}$ and the runs produce the same end model $\mathcal{Q}$ then the runs move the hull $H^{\mathcal{P},\Sigma}_{B}$ the same way. That is if i_1 and i_2 are B-iteration maps accoring to r_1 and r_2 respectively then $i_1 \restriction H^{\mathcal{P},\Sigma}_{B} = i_2 \restriction H^{\mathcal{P},\Sigma}_{B}$.*

Now we're ready to define our direct limit system. Let

$$\mathcal{F} = \{(P, \Sigma, \vec{B}) \quad | \quad \vec{B} \in \mathbb{B}(P^-, \Sigma)^{<\omega}, (P^-, \Sigma) \text{ satisfies Theorem 3.12}, (P, \Sigma) \text{ is } n\text{-suitable for some } n, \text{ and } (P, \Sigma) \text{ is strongly } \vec{B}\text{-iterable}\}.$$

The ordering on $\mathcal{F}$ is defined as follows:

$$\begin{aligned}(\mathcal{P},\Sigma,\vec{B}) \preccurlyeq (\mathcal{Q},\Lambda,\vec{C}) \quad iff \quad & \vec{B} \subseteq \vec{C}, \exists k \exists r (r \text{ is a run of } G^{\mathcal{P},\Sigma}_{B,k} \text{ with the last model } \mathcal{P}^* \\ & \text{such that } (\mathcal{P}^*)^- = \mathcal{Q}^-,\ \Sigma_{(\mathcal{P}^*)^-} = \Lambda, \mathcal{P}^* = \mathcal{Q}|(\eta^{+\omega})^{\mathcal{Q}} \\ & \text{ where } \mathcal{Q} \vDash \eta > o(Q^-) \ is\ Woodin).\end{aligned}$$

Suppose $(P,\Sigma,\vec{B}) \preccurlyeq (Q,\Lambda,\vec{C})$ then there is a unique map $\pi^{(\mathcal{P},\Sigma),(\mathcal{Q},\Delta)}_{\vec{B}} : H^{\mathcal{P},\Sigma}_{\vec{B}} \to H^{\mathcal{Q},\Lambda}_{\vec{B}}$. $(\mathcal{F},\preccurlyeq)$ is then directed. Let

$$\mathcal{M}_\infty = \text{direct limit of } (\mathcal{F},\preccurlyeq) \text{ under maps } \pi^{(\mathcal{P},\Sigma),(\mathcal{Q},\Delta)}_{\vec{B}}.$$

Also for each $(\mathcal{P},\Sigma,\vec{B}) \in \mathcal{F}$, let

$$\pi^{(\mathcal{P},\Sigma),\infty}_{\vec{B}} : H^{\mathcal{P},\Sigma}_{\vec{B}} \to \mathcal{M}_\infty$$

be the natural map.

Clearly, $\mathcal{M}_\infty \subseteq HOD$. But first, we need to show $\mathcal{F} \neq \emptyset$. In fact, we prove a stronger statement.

Theorem 3.16. *Suppose $(\mathcal{P},\Sigma)$ satisfies Theorem 3.12. Let $B \in \mathbb{B}(\mathcal{P},\Sigma)$. Then for each $1 \leq n < \omega$, there is a $\mathcal{Q}$ such that $\mathcal{Q}^-$ is a Σ-iterate of $\mathcal{P}^-$, $(Q,\Sigma_{\mathcal{Q}^-})$ is n-suitable and $(\mathcal{Q},\Sigma_{\mathcal{Q}^-},B) \in \mathcal{F}$.*

Proof. Suppose not. By Σ_1-reflection (Theorem 2.8), there is an transitive model N coded by a Suslin, co-Suslin set of reals such that $Code(\Sigma) \in \mathcal{P}(\mathbb{R})^N$ and

$$\begin{aligned} N \ \vDash \ & ZF^- + AD^+ + SMC + \text{“}\Theta \text{ exists and is successor in the Solovay sequence ”} + \\ & \text{“}\exists B \in \mathbb{B}(\mathcal{P},\Sigma)(\nexists \mathcal{Q}, n)((\mathcal{Q},\Sigma) \text{ is n-suitable and } (\mathcal{Q},\Sigma,B) \in \mathcal{F})\text{”}.\end{aligned}$$

We take a minimal such N and fix a $B \in \mathbb{B}(\mathcal{P},\Sigma)^N$ witnessing the failure of the Theorem in N. Using Theorem 2.7 and the assumption on N, there is an $x \in \mathbb{R}$ and a tuple $\langle N^*_x, \delta_x, \Sigma_x \rangle$ satisfying the conclusions of Theorem 2.7 relative to Γ- a good pointclass containing $(\mathcal{P}(\mathbb{R})^N, N's$ first order theory). Futhermore, let's assume that N^*_x Suslin captures $\langle A \mid A$ is projective in $\Sigma \rangle$). Let $\Omega = \mathcal{P}(\mathbb{R})^N$. For simplicity, we show that in N, there is a Σ-iterate $(\mathcal{R},\Sigma_{\mathcal{R}})$ such that there is a 1-suitable $(\mathcal{S},\Sigma_{\mathcal{R}})$ such that $(\mathcal{S},\Sigma_{\mathcal{R}},B) \in \mathcal{F}$.

By the assumption on N, $N \vDash V = K^\Sigma(\mathbb{R})$. Now N^*_x has club many $(\Sigma^2_1)^\Omega$ Woodins below δ_x by a standard argument (see [7]). Hence, the full background construction $L[E,\Sigma][\mathcal{P}]$ done in N^*_x will reach a model having ω Woodins (which are the first ω $(\Sigma^2_1)^\Omega$ Woodins in

N_x^*) and projecting across the sup of its first ω Woodins. Let $\mathcal{Q}$ be the first model in the construction with that property. By coring down if necessary, we may assume that $\mathcal{Q}$ is sound. Let $\langle \delta_i^{\mathcal{Q}} \mid i < \omega \rangle$ be the first ω Woodins of $\mathcal{Q}$ above $o(\mathcal{P})$. A similar self-explanatory notation will be used to denote the Woodins of any Λ-iterate of $\mathcal{Q}$. Hence $\rho_\omega(\mathcal{Q}) < sup_{i<\omega}\delta_i$. Let Λ (which extends Σ) be the strategy of $\mathcal{Q}$ induced from the background universe. Λ is Ω-fullness preserving. At this point it's not clear that Λ has branch condensation. The proof of Theorem 2.11 doesn't generalize as it's not clear what the corresponding notion of a self-justifying-system for sets in $\mathbb{B}(\mathcal{P}, \Sigma)$ is.

We in fact show a bit more. We show that an iterate $(\mathcal{R}, \Lambda_{\mathcal{R}})$ of $(\mathcal{Q}, \Lambda)$ has strong B condensation in that if $i : \mathcal{R} \to \mathcal{S}$ is according to $\Lambda_{\mathcal{R}}$ and below $\delta_0^{\mathcal{Q}}$ and $j : \mathcal{R} \to \mathcal{W}$ is such that there is a $k : \mathcal{W} \to \mathcal{S}$ such that $i = k \circ j$ then $i(\tau^{\mathcal{R}}_{B,\delta_0^{\mathcal{R}}}) = \tau^{\mathcal{S}}_{B,\delta_0^{\mathcal{S}}}$, $\mathcal{W}$ is full, and $k^{-1}(\tau^{\mathcal{S}}_{B,\delta_0^{\mathcal{S}}}) = \tau^{\mathcal{W}}_{B,j(\delta_0^{\mathcal{R}})}$. That we get $\mathcal{W}$ being full is easy because $\Lambda \notin \mathcal{N}$. So we only need to prove the other two clauses. We also get strong B-iterability by the Dodd-Jensen property of $\Lambda_{\mathcal{R}}$. Once we have this pair $(\mathcal{R}, \Lambda_{\mathcal{R}})$, we can just let our desired $\mathcal{S}$ to be $\mathcal{R}|((\delta_0^{\mathcal{R}})^{+\omega})^{\mathcal{R}}$. Suppose not. Using the property of $\mathcal{Q}$ and the relativized (to Σ) Prikry forcing in N (see [11]), we get that for any n, there is an iterate $\mathcal{R}$ of $\mathcal{Q}$ (above $\delta_0^{\mathcal{Q}}$) extending a Prikry generic and having N as the (new) derived model (computed at the sup of the first ω Woodins above $o(\mathcal{P})$). Furthermore, this property holds for any Λ iterate of $\mathcal{Q}$. Without going further into details of the techniques used in [11], we remark that if $\mathcal{R}$ is an $\mathbb{R}$-genericity iterate of $\mathcal{Q}$, then the new derived model of $\mathcal{R}$ is N. In other words, once we know one such $\mathbb{R}$-genericity iterate of $\mathcal{Q}$ realizes N as its derived model then all $\mathbb{R}$-genericity iterates of $\mathcal{Q}$ do. Let (ϕ, s) define B over N, i.e.

$$(\mathcal{R}, \Psi, x, y) \in B \ iff \ N \vDash \phi[((\mathcal{R}, \Psi, x, y)), s].$$

From now to the end of the proof, all stacks on $\mathcal{Q}$ or its iterates thereof are below the $\delta_0^{\mathcal{Q}}$ or its image. By our assumption, there is $\langle \vec{\mathcal{T}}_i, \vec{\mathcal{S}}_i, \mathcal{Q}_i, \mathcal{R}_i, \pi_i, \sigma_i, j_i \mid i < \omega \rangle \in N$ such that

1. $\mathcal{Q}_0 = \mathcal{Q}$; $\vec{\mathcal{T}}_0$ is a stack on $\mathcal{Q}$ according to Λ with last model $\mathcal{Q}_1$; $\pi_0 = i^{\vec{\mathcal{T}}_0}$; $\vec{\mathcal{S}}_0$ is a stack on $\mathcal{Q}$ with last model $\mathcal{R}_0$; $\sigma_0 = i^{\vec{\mathcal{S}}_0}$; and $j_0 : \mathcal{R}_0 \to \mathcal{Q}_1$.

2. $\vec{\mathcal{T}}_i$ is a stack on $\mathcal{Q}_i$ according to Λ with last model $\mathcal{Q}_{i+1}$; $\pi_i = i^{\vec{\mathcal{T}}_i}$; $\vec{\mathcal{S}}_i$ is a stack on $\mathcal{Q}_i$ with last model $\mathcal{R}_i$; $\sigma_i = i^{\vec{\mathcal{S}}_i}$; $j_0 : \mathcal{R}_i \to \mathcal{Q}_{i+1}$.

3. for all k, $\pi_k = j_k \circ \sigma_k$.

4. for all k, $\pi_k(\tau^{\mathcal{Q}_k}_{B,\delta_0^{\mathcal{Q}_k}}) \neq \tau^{\mathcal{Q}_{k+1}}_{B,\delta_0^{\mathcal{Q}_{k+1}}}$ or $j_k(\tau^{\mathcal{R}_k}_{B,\delta_0^{\mathcal{R}_k}}) \neq \tau^{\mathcal{Q}_{k+1}}_{B,\delta_0^{\mathcal{Q}_{k+1}}}$.

Let $\mathcal{Q}_\omega$ be the direct limit of the $\mathcal{Q}_i$'s under maps π_i's. First we rename the $\langle \mathcal{Q}_i, \mathcal{R}_i, \pi_i, \sigma_i, j_i \mid i < \omega\rangle$ into $\langle \mathcal{Q}_i^0, \mathcal{R}_i^0, \pi_i^0, \sigma_i^0, j_i^0 \mid i < \omega\rangle$. We then assume that N is countable (by working with a countable elementary substructure of N) and fix (in V) $\langle x_i \mid i < \omega\rangle$- a generic enumeration of $\mathbb{R}$. Using our assumption on $\mathcal{Q}$, we get $\langle \mathcal{Q}_i^n, \mathcal{R}_i^n, \pi_i^n, \sigma_i^n, j_i^n, \tau_1^n, k_i^n \mid n, i \leq \omega\rangle$ such that

1. $\mathcal{Q}_i^\omega$ is the direct limit of the $\mathcal{Q}_i^n$'s under maps τ_i^n's for all $i \leq \omega$.
2. $\mathcal{R}_i^\omega$ is the direct limit of the $\mathcal{R}_i^n$'s under maps k_i^n's for all $i < \omega$.
3. $\mathcal{Q}_\omega^n$ is the direct limit of the $\mathcal{Q}_i^n$'s under maps π_i^n's.
4. for all $n \leq \omega$, $i < \omega$, $\pi_i^n : \mathcal{Q}_i^n \to \mathcal{Q}_{i+1}^n$; $\sigma_i^n : \mathcal{Q}_i^n \to \mathcal{R}_i^n$; $j_i^n : \mathcal{R}_i^n \to \mathcal{Q}_{i+1}^n$ and $\pi_i^n = j_i^n \circ \sigma_i^n$.
5. Derived model of the $\mathcal{Q}_i^n$'s, $\mathcal{R}_i^n$'s is N.

Then we start by iterating $\mathcal{Q}_0^0$ above $\delta_0^{\mathcal{Q}_0^0}$ to $\mathcal{Q}_0^1$ to make x_0-generic at $\delta_1^{\mathcal{Q}_0^1}$. During this process, we lift the genericity iteration tree to all $\mathcal{R}_n^0$ for $n < \omega$ and $\mathcal{Q}_n^0$ for $n \leq \omega$. We pick branches for the tree on $\mathcal{Q}_0^0$ by picking branches for the lift-up tree on $\mathcal{Q}_\omega^0$ using $\Lambda_{\mathcal{Q}_\omega^0}$. Let $\tau_0^0 : \mathcal{Q}_0^0 \to \mathcal{Q}_0^1$ be the iteration map and $\mathcal{W}$ be the end model of the lift-up tree on $\mathcal{Q}_\omega^0$. We then iterate the end model of the lifted tree on $\mathcal{R}_0^0$ to $\mathcal{R}_0^1$ to make x_0 generic at $\delta_1^{\mathcal{R}_0^1}$ with branches being picked by lifting the iteration tree onto $\mathcal{W}$ and using the branches according to $\Lambda_{\mathcal{W}}$. Let $k_0^0 : \mathcal{R}_0^0 \to \mathcal{R}_0^1$ be the iteration embedding, $\sigma_0^1 : \mathcal{Q}_0^1 \to R_0^1$ be the natural map, and $\mathcal{X}$ be the end model of the lifted tree on the $\mathcal{W}$ side. We then iterate the end model of the lifted stack on $\mathcal{Q}_1^0$ to $\mathcal{Q}_1^1$ to make x_0 generic at $\delta_1^{\mathcal{Q}_1^1}$ with branches being picked by lifting the tree to $\mathcal{X}$ and using branches picked by $\Lambda_{\mathcal{X}}$. Let $\tau_1^0 : \mathcal{Q}_1^0 \to \mathcal{Q}_1^1$ be the iteration embedding, $j_0^1 : \mathcal{R}_0^1 \to \mathcal{Q}_1^1$ be the natural map, and $\pi_0^1 = j_0^1 \circ \sigma_0^1$. Continue this process of making x_0 generic for the later models $\mathcal{R}_n^0$'s and $\mathcal{Q}_n^0$'s for $n < \omega$. We then let $\mathcal{Q}_\omega^1$ be the direct limit of the $\mathcal{Q}_n^1$ under maps π_n^1's. We then start at $\mathcal{Q}_0^1$ and repeat the above process to make x_1 generic appropriate iterates of $\delta_2^{\mathcal{Q}_0^1}$ etc. This whole process define models and maps $\langle \mathcal{Q}_i^n, \mathcal{R}_i^n, \pi_i^n, \sigma_i^n, j_i^n, \tau_1^n, k_i^n \mid n, i \leq \omega\rangle$ as described above. See Figure 2.

Note that by our construction, for all $n < \omega$, the maps π_n^0's and τ_ω^n's are via Λ or its appropriate tails; furthermore, $\mathcal{Q}_\omega^\omega$ is wellfounded and full (with respect to mice in N). This in turns implies that the direct limits $\mathcal{Q}_n^\omega$'s and $\mathcal{R}_n^\omega$'s are wellfounded and full. We must then have that for some k, for all $n \geq k$, $\pi_n^\omega(s) = s$. This implies that for all $n \geq k$

$$\pi_n^\omega(\tau^{\mathcal{Q}_n^\omega}_{B,\delta_0^{\mathcal{Q}_n^\omega}}) = \tau^{\mathcal{Q}_{n+1}^\omega}_{B,\delta_0^{\mathcal{Q}_{n+1}^\omega}}.$$

We can also assume that for all $n \geq k$, $\sigma_n^\omega(s) = s, j_n^\omega(s) = s$. Hence

$$\sigma_n^\omega(\tau^{\mathcal{Q}_n^\omega}_{B,\delta_0^{\mathcal{Q}_n^\omega}}) = \tau^{\mathcal{R}_n^\omega}_{B,\delta_0^{\mathcal{R}_n^\omega}}.;$$

$$\begin{tikzcd}
\mathcal{Q}^0_0 \arrow[rr, "\pi^0_0"] \arrow[dr, "\sigma^0_0"'] \arrow[dd, "\tau^0_0"'] & & \mathcal{Q}^0_1 \arrow[rr, "\pi^0_1"] \arrow[dr, "\sigma^0_1"'] \arrow[dd, "\tau^0_1"'] & & \mathcal{Q}^0_2 \arrow[r, "\pi^0_2"] \arrow[dd, "\tau^0_2"'] & \cdots & \mathcal{Q}^0_\omega \arrow[dd, "\tau^0_\omega"'] \\
& \mathcal{R}^0_0 \arrow[ur, "j^0_0"'] \arrow[dd, "k^0_0"] & & \mathcal{R}^0_1 \arrow[ur, "j^0_1"'] \arrow[dd, "k^0_1"] & & & \\
\mathcal{Q}^1_0 \arrow[rr, "\pi^1_0"] \arrow[dr, "\sigma^1_0"'] \arrow[dd, "\tau^1_0"'] & & \mathcal{Q}^1_1 \arrow[rr, "\pi^1_1"] \arrow[dr, "\sigma^1_1"'] \arrow[dd, "\tau^1_1"'] & & \mathcal{Q}^1_2 \arrow[r, "\pi^1_2"] \arrow[dd, "\tau^1_2"'] & \cdots & \mathcal{Q}^1_\omega \arrow[dd, "\tau^1_\omega"'] \\
& \mathcal{R}^1_0 \arrow[ur, "j^1_0"'] \arrow[dd, "k^1_0"] & & \mathcal{R}^1_1 \arrow[ur, "j^1_1"'] \arrow[dd, "k^1_1"] & & & \\
\mathcal{Q}^2_0 \arrow[rr, "\pi^2_0"] \arrow[dr, "\sigma^2_0"'] \arrow[dd, "\tau^2_0"'] & & \mathcal{Q}^2_1 \arrow[rr, "\pi^2_1"] \arrow[dr, "\sigma^2_1"'] \arrow[dd, "\tau^2_1"'] & & \mathcal{Q}^2_2 \arrow[r, "\pi^2_2"] \arrow[dd, "\tau^2_2"'] & \cdots & \mathcal{Q}^2_\omega \arrow[dd, "\tau^2_\omega"'] \\
& \mathcal{R}^2_0 \arrow[ur, "j^2_0"'] \arrow[dd, "k^2_0"] & & \mathcal{R}^2_1 \arrow[ur, dash] \arrow[dd, "k^2_1"] & & & \\
\vdots & & \vdots & & \vdots & & \vdots \\
& \vdots & & \vdots & & & \\
\mathcal{Q}^\omega_0 \arrow[rr, "\pi^\omega_0"] \arrow[dr, "\sigma^\omega_0"'] & & \mathcal{Q}^\omega_1 \arrow[rr, "\pi^\omega_1"] \arrow[dr, "\sigma^\omega_1"'] & & \mathcal{Q}^\omega_2 \arrow[r, "\pi^\omega_2"] & \cdots & \mathcal{Q}^\omega_\omega \\
& \mathcal{R}^\omega_0 \arrow[ur, "j^\omega_0"'] & & \mathcal{R}^\omega_1 \arrow[ur, "j^\omega_1"'] & & &
\end{tikzcd}$$

Figure 2: The process in Theorem 3.16

$$j_n^\omega(\tau^{\mathcal{R}_n^\omega}_{B,\delta_0^{\mathcal{R}_n^\omega}}) = \tau^{\mathcal{Q}^\omega_{n+1}}_{B,\delta_0^{\mathcal{Q}^\omega_{n+1}}}.;$$

This is a contradiction, hence we're done. □

Remark 3.17. The proof of Theorem 3.16 also shows that if $(\mathcal{P}, \Sigma)$ is n-suitable and $(\mathcal{P}, \Sigma, B) \in \mathcal{F}$ and $C \in \mathbb{B}(\mathcal{P}^-, \Sigma)$ then there is a B-iterate $\mathcal{Q}$ of $\mathcal{P}$ such that $(\mathcal{Q}, \Sigma_{\mathcal{Q}^-}, B \oplus C) \in \mathcal{F}$; in fact, $\mathcal{Q}$ has strong $B \oplus C$-condensation as defined in the proof of Theorem 3.16.

It is easy to see that $\mathcal{M}_\infty | \theta_\alpha = V^{HOD}_{\theta_\alpha}$. Let $\langle \eta_i \mid i < \omega \rangle$ be the increasing enumeration of Woodin cardinals in $\mathcal{M}_\infty$ larger than θ_α. Theorem 3.16 is used to show that $\mathcal{M}_\infty$ is large enough in that

Lemma 3.18. *1. $\mathcal{M}_\infty$ is well-founded.*

2. $\mathcal{M}_\infty | \eta_0 = V^{HOD}_\Theta$. In particular, $\eta_0 = \Theta$.

Proof. We prove (1) and (2) simultaneously. Toward a contradiction, suppose not. By Σ_1-reflection (Theorem 2.8), there is a transitive model N coded by a Suslin, co-Suslin set of reals such that $Code(\Sigma) \in \mathcal{P}(\mathbb{R})^N$ and

$$\begin{aligned} N \ \models\ & ZF^- + DC + SMC + \text{“}\Theta \text{ exists and is successor in the Solovay sequence ”} + \\ & \text{“(1) and (2) do not both hold”}. \end{aligned}$$

We take a minimal such N and let $\Omega = \mathcal{P}(\mathbb{R})^N$. We get $N \models V = K^{\Sigma}(\mathbb{R})$ and a $(\mathcal{Q}, \Lambda)$ with the property that for all $B \in \mathbb{B}(P, \Sigma)^N$, there is a Λ iterate $\mathcal{R}$ of $\mathcal{Q}$ that strongly respects B. $(\mathcal{Q}, \Lambda)$ also has the property that any Λ iterate $\mathcal{R}$ of $\mathcal{Q}$ can be further iterated by $\Lambda_{\mathcal{R}}$ to $\mathcal{S}$ such that N is the derived model of $\mathcal{S}$.

Fix $\langle \alpha_i \mid i < \omega \rangle$ a cofinal in Θ^{Ω} sequence of ordinals. Such a sequence exists since $\Omega = Env((\Sigma^2_1)^N)$. For each n, let

$$\begin{aligned} D_n = \{(\mathcal{R}, \Psi, x, y) \quad \mid \quad & (\mathcal{R}, \Psi) \text{ is a hod pair equivalent to } (\mathcal{P}, \Sigma),\ x \text{ codes } \mathcal{R}, \\ & y \in \text{ the least } OD^N_{\Psi} \text{ set of reals with Wadge rank } \geq \alpha_n\} \end{aligned}$$

Clearly, for all n, $D_n \in \mathbb{B}(\mathcal{P}, \Sigma)^N$. Without loss of generality, we may assume Λ strongly respects all the D_n's and the derived model of $\mathcal{Q}$ is N. Let $\vec{D} = \langle D_n \mid n < \omega \rangle$. Before proving the next claim, let us introduce the following notion. First let for a set $A(A \subseteq \mathbb{R}$ or $A \in \mathbb{B}(\mathcal{P}, \Sigma))$, $\tau^{Q,0}_{A,m}$ be the canonical capturing term for A in $\mathcal{Q}$ at $(\delta^{+m}_0)^{\mathcal{Q}}$. Set

$$\begin{aligned} \gamma^{\mathcal{Q},0}_{D_i,m} &= sup\{H^{\mathcal{Q}}_1(P \cup \{\tau^{\mathcal{Q},0}_{D_i,m}\}) \cap \delta_0\}; \\ \gamma^{\mathcal{Q},0}_{D_i} &= sup_{m<\omega} \gamma^{\mathcal{Q},0}_{D_i,m}. \end{aligned}$$

Claim 1. *For any Λ-iterate $(\mathcal{S}, \Upsilon)$ of $\mathcal{Q}$. Suppose $i : \mathcal{Q} \to \mathcal{S}$ is the itaration map. Then*

$$i(\delta_0) = sup_{i<\omega} \gamma^{\mathcal{S},0}_{D_i}.$$

Proof. Working in N, let $\langle A_i \mid i < \omega \rangle$ be a sequence of OD_{Σ} sets such that A_0 is a universal $\Sigma^2_1(\Sigma)$ set; $A_1 = \mathbb{R} \backslash A_0$; the $\langle A_i \mid i \geq 2 \rangle$ is a $\Pi^2_1(\Sigma)$-semiscale on A_1. Suppose ϕ_i and $s_i \in ORD^{<\omega}$ are such that

$$x \in A_i \ iff \ N \models \phi_i[\Sigma, s_i, x]$$

Now for each i, let

$$\begin{aligned} A_i^* = \{(\mathcal{R}, \Psi, x, y) \quad | \quad & (\mathcal{R}, \Psi) \text{ is a hod pair equivalent to } (\mathcal{P}, \Sigma),\ x \text{ codes } \mathcal{R}, \\ & N \vDash \phi_i[\Psi, s_i, y]\} \end{aligned}$$

Aside from the assumption about $(\mathcal{Q}, \Lambda)$ above ,we also assume Λ is guided by $\langle A_i \mid i < \omega\rangle$ for stacks above $\mathcal{P}$ and below δ_0. This is possible by relativizing to Σ the proof of a similar fact in the case $\Theta = \theta_0$. This means

$$\delta_0 = sup_{i<\omega}\gamma^{\mathcal{Q},0}_{A_i^*}.$$

This fact in turns implies

$$\delta_0 = sup_{i<\omega}\gamma^{\mathcal{Q},0}_{D_i}.$$

To see this, fix an A_i^*. We'll show that there is a j such that $\gamma^{\mathcal{Q},0}_{D_j} \geq \gamma^{\mathcal{Q},0}_{A_i^*}$. Well, fix a real coding $\mathcal{P}$ and let j be such that

$$w(A_i) = w((A_i^*)_{(\mathcal{P},\Sigma,x)}) \leq w((D_j)_{(\mathcal{P},\Sigma,x)}).$$

Let z be a real witnessing the reduction. Then there is a map $i : \mathcal{Q} \to \mathcal{R}$ such that

1. i is according to Λ and the iteration is above $\mathcal{Q}^- = \mathcal{P}$;

2. z is generic for the extender algebra $\mathbb{A}$ of $\mathcal{R}$ at $\delta^{\mathcal{R}}$.

Note that $i(\tau^{\mathcal{Q}}_{A_i^*}) = \tau^{R}_{A_i^*}$, $i(\tau^{\mathcal{Q}}_{D_j}) = \tau^{\mathcal{R}}_{D_j}$, and $\mathcal{R}[z] \vDash \tau_{A_i^*} \leq_w \tau_{D_j}$ via z. Hence $\tau^{\mathcal{R}}_{A_i^*} \in X = \{\tau \in \mathcal{R}^{\mathbb{A}} \mid (\exists p \in \mathbb{A})(p \Vdash_R \tau \leq_w \tau_{D_j} \text{ via } \dot{z})\}$ and $|X|^{\mathcal{R}} < \delta^{\mathcal{R}}$ (by the fact that the extender algebra $\mathbb{A}$ is $\delta^{\mathcal{R}}$-cc). But X is definable over $\mathcal{R}$ from $\tau^{R}_{D_j}$, hence $|X|^{\mathcal{R}} < \gamma^{R,0}_{D_j}$. Since $\tau^{R}_{A_i^*} \in X$, $\gamma^{\mathcal{R},0}_{A_i^*} \leq \gamma^{\mathcal{R},0}_{D_j}$ which in turns implies $\gamma^{\mathcal{Q},0}_{A_i^*} \leq \gamma^{\mathcal{Q},0}_{D_j}$.

Now to finish the claim, let $(\mathcal{S}, \Upsilon)$ be a Λ iterate of $\mathcal{Q}$. Suppose $i : \mathcal{Q} \to \mathcal{S}$ is the iteration map. Let $\mathcal{R} = i(\mathcal{P})$ and $\Sigma_{\mathcal{Q}}$ be the tail of Σ under the iteration. We claim that

$$i(\delta_0) = sup_{i<\omega}\gamma^{S,0}_{D_i}.(*)$$

This is easily seen to finish the proof of Claim 1. To see $(*)$, we repeat the proof of the previous part applied to $(\mathcal{S}, \Upsilon)$ and $\langle B_i \mid i < \omega\rangle$ where B_0 is a universal $\Sigma^2_1(\Sigma_{\mathcal{Q}})$; $B_1 = \mathbb{R}\backslash B_0$; $\langle B_i \mid i \geq 2\rangle$ is a $\Pi^2_1(\Sigma_{\mathcal{Q}})$-semiscale on B_1. We may assume $(\mathcal{S}, \Upsilon)$ is guided by $\langle B_i \mid i < \omega\rangle$ for stacks above R and below $i(\delta_0)$. Now we are in the position to apply the exact same argument as above and conclude that $(*)$ holds. Hence we're done. $\square$

The proof of claim 1 shows Λ has branch condensation, hence the direct limit $\mathcal{M}_\infty(\mathcal{Q},\Lambda)$ is defined and is wellfounded. This implies that in N, $\mathcal{M}_\infty$ is wellfounded. Let $\langle \delta_i \mid i<\omega\rangle$ be the first ω Woodins of $\mathcal{Q}$ above $\mathcal{Q}^-$ and $i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\infty} : \mathcal{Q} \to \mathcal{M}_\infty(\mathcal{Q},\Lambda)$ be the iteration embedding according to Λ and $\langle \eta_n \mid n<\omega\rangle = \langle i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\infty}(\delta_i) \mid i<\omega\rangle$. For $(\mathcal{R},\Lambda_{\mathcal{R}})$ and iterate of $(\mathcal{Q},\Lambda)$, let $i^{\mathcal{R},\Lambda_{\mathcal{R}}}_{\mathcal{R},\infty}$ have the obvious meaning and $i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{R}}$ be the iteration map according to Λ. Note that in N, $\mathcal{M}_\infty(\mathcal{Q},\Lambda)|\eta_n = \mathcal{M}_\infty|\eta_n$ for all n.

Claim 2. $\eta_0 = \Theta^\Omega$.

Proof. Working in N, we first claim that

$$\mathcal{M}_\infty(\mathcal{Q},\Lambda)|\eta_0 = V^{HOD}_{\eta_0}. \quad (*)$$

To show $(*)$, it is enough to show that if $A \subseteq \alpha < \eta_0$ and A is OD then $A \in \mathcal{M}_\infty(\mathcal{Q},\Lambda)$. To see this, let i be such that $\gamma^{\mathcal{M}_\infty(\mathcal{Q},\Lambda),0}_{D_i} > \alpha$ (such an i exists by the proof of Claim 1). Let

$$\begin{array}{lcl} C = \{(\mathcal{R},\Psi,x,y) & \mid & (\mathcal{R},\Psi) \text{ is a hod pair equivalent to } (\mathcal{P},\Sigma),\ x \text{ codes } \mathcal{R},\ y \text{ codes } (N,\gamma) \\ & & \text{such that } (\mathcal{N},\Psi) \text{ is 1-suitable, } \mathcal{N} \text{ is strongly } D_i \text{ iterable via a} \\ & & \text{quasi-strategy } \Phi \text{ extending } \Psi,\ \gamma < \gamma^{\mathcal{N},0}_{D_i},\ \pi^{(\mathcal{N},\Psi),\infty}_{D_i}(\gamma) \in A\}. \end{array}$$

By replacing $\mathcal{Q}$ by an iterate we may assume $(\mathcal{Q},\Lambda)$ is C-iterable. Let $\tau^{\mathcal{Q}}_C = \tau^{\mathcal{Q}}_{C,(\delta_0^{+\omega})^{\mathcal{Q}}}$ and $\tau_C = i^{(\mathcal{Q},\Lambda)}_{\mathcal{Q},\infty}(\tau^{\mathcal{Q}}_C)$. The following equivalence is easily shown by a standard computation:

$$\begin{array}{ll} \xi \in A \ iff \ \mathcal{M}_\infty(\mathcal{Q},\Lambda) \vDash\!\Vdash_{Col(\omega,\eta_0^{+\omega})} & \text{“if } x \text{ codes } i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\infty}(\mathcal{P}), y \text{ codes } (\mathcal{M}_\infty(\mathcal{Q},\Lambda)|\eta_0^{+\omega},\xi) \\ & \quad \text{then } (x,y) \in \tau_C\text{”.} \end{array}$$

For the reader's convenience, we'll show why the above equivalence holds. First suppose $\xi \in A$. Let $(\mathcal{S},\Xi) \in I(\mathcal{Q},\Lambda)$ be such that there is a $\gamma < \gamma^{\mathcal{S},0}_{D_i}$ and $i^{\mathcal{S},\Xi}_{\mathcal{S},\infty}(\gamma) = \xi$. Then we have (letting $\nu = i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\delta_0)$)

$$\mathcal{S} \vDash\!\Vdash_{Col(\omega,\nu^{+\omega})} \text{“if } x \text{ codes } i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\mathcal{P}), y \text{ codes } (S|\nu^{+\omega},\gamma) \text{ then } (x,y) \in i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\tau^{\mathcal{Q},\Lambda}_C)\text{”.}$$

By applying $i^{\mathcal{S},\Xi}_{\mathcal{S},\infty}$ to this ,we get

$$\mathcal{M}_\infty(\mathcal{Q},\Lambda) \vDash\!\Vdash_{Col(\omega,\eta_0^{+\omega})} \text{“if } x \text{ codes } i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\infty}(\mathcal{P}), y \text{ codes } (\mathcal{M}_\infty(\mathcal{Q},\Lambda)|\eta_0^{+\omega},\xi) \text{ then } (x,y) \in \tau_C\text{”.}$$

Now to show $(\Leftarrow)$, let $(\mathcal{S},\Xi) \in I(\mathcal{Q},\Lambda)$ be such that for some $\gamma < \gamma^{\mathcal{S},0}_{D_i}$, $\xi = i^{\mathcal{S},\Xi}_{\mathcal{S},\infty}(\gamma)$. Let

$\nu = i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\delta_0)$, we have

$$\mathcal{S} \vDash\Vdash_{Col(\omega,\nu^{+\omega})} \text{“if } x \text{ codes } i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\mathcal{P}), y \text{ codes } (\mathcal{S}|\nu^{+\omega},\gamma) \text{ then } (x,y) \in i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\tau^{\mathcal{Q},\Lambda}_{C})\text{”}.$$

This means there is a quasi-strategy Ψ on $\mathcal{S}(0)$ ($\mathcal{S}(0) = \mathcal{S}|(\nu^{+\omega})^{\mathcal{S}}$) such that $(\mathcal{S}(0), i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\Sigma))$ is 1-suitable, Ψ extends $i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\Sigma))$, and Ψ is D_i-iterable. We need to see that $\pi^{(\mathcal{S}(0),i^{\mathcal{Q},\Lambda}_{\mathcal{Q},\mathcal{S}}(\Sigma)),\infty}_{D_i}(\gamma) = \xi$. But this is true by the choice of D_i, $\xi = i^{\mathcal{S},\Xi}_{S,\infty}(\gamma)$, and the fact that Ψ agrees with Ξ on how ordinals below $\gamma^{\mathcal{S},0}_{D_i}$ are mapped.

The equivalence above shows $A \in \mathcal{M}_\infty(Q,\Lambda)$, hence completes the proof of $(*)$. $(*)$ in turns shows that η_0 is a cardinal in HOD and $\eta_0 \leq \Theta$ (otherwise, $HOD|\eta_0 = \mathcal{M}_\infty(Q,\Lambda)|\eta_0 \vDash \Theta$ is not Woodin while $HOD \vDash \Theta$ is Woodin).

Next, we show expectedly that

$$\eta_0 = \Theta. \quad (**)$$

Suppose toward a contradiction that $\eta_0 < \Theta$. Let $\mathcal{Q}(0) = \mathcal{Q}|(\delta_0^{+\omega})^{\mathcal{Q}}$, $\Lambda_0 = \Lambda|\mathcal{Q}(0)$, and $\mathcal{M}_\infty(\mathcal{Q},\Lambda)(0) = \mathcal{M}_\infty(\mathcal{Q},\Lambda)|(\eta_0^{+\omega})^{\mathcal{M}_\infty(\mathcal{Q},\Lambda)}$. Let $\pi = i \restriction \mathcal{Q}(0)$; so π is according to Λ_0. By the Coding Lemma and our assumption that $\eta_0 < \Theta$, $\pi, \mathcal{M}_\infty(Q,\Lambda)(0) \in N$. From this, we can show $\Lambda_0 \in N$ by the following computation: $\Lambda_0(\vec{\mathcal{T}}) = b$ if and only if

1. the part of $\vec{\mathcal{T}}$ based on P is according to Σ;

2. if $i^{\vec{\mathcal{T}}}_b$ exists then there is a $\sigma : \mathcal{M}^{\vec{\mathcal{T}}}_b \to \mathcal{M}_\infty(Q,\Lambda)(0)$ such that $\pi = \sigma \circ i^{\vec{\mathcal{T}}}_b$;

3. $\vec{\mathcal{T}}^\frown\mathcal{M}^{\vec{\mathcal{T}}}_b$ is Q-structure guided.

By branch condensation of Λ_0, (1),(2), and (3) indeed define Λ_0 in N. This means Λ_0 is OD^N from Σ (and some real x); hence $\Lambda_0 \in N$. So suppose $\gamma = w(Code(\Lambda_0)) < \Theta^\Omega$. In N, let

$$\begin{aligned} B = \{(\mathcal{R},\Psi,x,y) \quad | \quad & (\mathcal{R},\Psi) \text{ is a hod pair equivalent to } (\mathcal{P},\Sigma),\ x \text{ codes } \mathcal{R},\ y \in A_{\mathcal{R}} \\ & \text{where } A_{\mathcal{R}} \text{ is the least } OD(Code(\Psi)) \text{ set such that } w(A_{\mathcal{R}}) > \gamma\} \end{aligned}$$

Then $B \in \mathbb{B}(\mathcal{P},\Sigma)^N$. We may assume Λ_0 respects B. It is then easy to see that whenever $(\mathcal{R},\Lambda_{\mathcal{R}}) \in I(Q(0),\Lambda_0)$ (also let $\mathcal{S} \lhd \mathcal{R}$ be the iterate of $\mathcal{P}$), $w(Code(\Lambda_{\mathcal{R}})) \geq w(A_{\mathcal{R}})$ because $\Lambda_{\mathcal{R}}$ can compute membership of $A_{\mathcal{R}}$ by performing genericity iterations (above $\mathcal{S}$) to make reals generic. This means $w(Code(\Lambda_{\mathcal{R}})) > \gamma = w(Code(\Lambda_0))$. This contradicts the fact that $w(Code(\Lambda_{\mathcal{R}})) = w(Code(\Lambda_0))$. □

The proof of Claim 1 and Claim 2 shows that the fragment of Λ on stacks above P and below δ_0 is guided by $\langle D_i \mid i < \omega \rangle$. $\square$

Now we define a strategy Σ_∞ for $\mathcal{M}_\infty$ extending the strategy Σ_∞^- of $\mathcal{M}_\infty^- = V_{\theta_\alpha}^{HOD}$. Let $(\mathcal{P}, \Sigma, A) \in \mathcal{F}$ and suppose $\mathcal{P}$ is n-suitable with $\langle \delta_i \mid i < n \rangle$ being the sequence of Woodins of $\mathcal{P}$ above $\mathcal{P}^-$, let $\tau_{A,k}^{\mathcal{M}_\infty}$ = common value of $\pi_{\vec{B},\infty}^{\mathcal{P},\Sigma}(\tau_{A,\delta_k}^{\mathcal{P}})$. Σ_∞ will be defined (in V) for trees on $\mathcal{M}_\infty|\eta_0$ in $\mathcal{M}_\infty$. For $k \geq n$, $\mathcal{M}_\infty \vDash$ "$Col(\omega, \eta_n) \times Col(\omega, \eta_k) \Vdash (\tau_{A,n}^{\mathcal{M}_\infty})_g = (\tau_{A,k}^{M_\infty})_h \cap \mathcal{M}_\infty[g]$" where g is $Col(\omega, \eta_n)$ generic and h is $Col(\omega, \eta_k)$ generic and $(\tau_{A,n}^{\mathcal{M}_\infty})_g$ is understood to be $A_{(\mathcal{M}_\infty^-, \Sigma_\infty^-)} \cap \mathcal{M}_\infty[g]$. This is just saying that the terms cohere with one another.

Let $\lambda^{\mathcal{M}_\infty} = sup_{i<\omega} \eta_i$. Let G be $Col(\omega, \lambda^{\mathcal{M}_\infty})$ generic over $\mathcal{M}_\infty$. Then $\mathbb{R}_G^*$ is the symmetric reals and $A_G^* := \cup_k (\tau_{A,k}^{\mathcal{M}_\infty})_{G|\eta_k}$.

Proposition 3.19. *For all $A \in \mathbb{B}(\mathcal{M}_\infty^-, \Sigma_\infty^-)$, $L(A_G^*, \mathbb{R}_G^*) \vDash AD^+$*

Proof. We briefly sketch the proof of this since the techniques involved have been fully spelled out in Section 2. If not, reflect the situation down to a model N coded by a Suslin co-Suslin set. Next get a "next mouse" $\mathcal{N}$ with ω Woodin cardinals that iterates out to (possibly a longer mouse than) $\mathcal{M}_\infty^N$. We can and do assume that an iterate of $\mathcal{N}$ has derived model is $K^\Sigma(\mathbb{R})$ of N, where $(\mathcal{P}, \Sigma)$ is a hod pair giving us $HOD|\theta_\alpha$; the proof of this fact uses a relative-to-Σ Prikry forcing (see [11]) and S-constructions (see [3]). From now on, we work inside the reflected universe N.

Let $A \subseteq \mathbb{B}(\mathcal{M}_\infty^-, \Sigma_\infty^-)$ be the least OD set such that $L(A_G^*, \mathbb{R}_G^*) \nvDash AD^+$. Then there is an iterate $\mathcal{M}$ of $\mathcal{N}$ having preimages of all the terms $\tau_{A,k}^{M_\infty}$. We may assume $\mathcal{M}$ has derived model $K^\Sigma(\mathbb{R})$. Since we have AD^+, $\mathcal{M}$ thinks that its derived model (in this case is $K^\Sigma(\mathbb{R})$) satisfies that $L(A_{(\mathcal{P},\Sigma)}, \mathbb{R}) \vDash AD^+$, where we reuse $(\mathcal{P}, \Sigma)$ for an equivalent (but possibly different) hod pair from the original one. Now iterate $\mathcal{M}$ to $\mathcal{Q}$ such that M_∞ is a proper initial segment of $\mathcal{Q}$. By elementarity $L(A_G^*, \mathbb{R}_G^*) \vDash AD^+$. This is a contradiction. $\square$

Definition 3.20. *Given a normal tree $\mathcal{T} \in \mathcal{M}_\infty$ and $\mathcal{T}$ is based on $\mathcal{M}_\infty|\theta_0$. $\mathcal{T}$ is by Σ_∞ if the following hold (the definition is similar for finite stacks):*

- *If $\mathcal{T}$ is short then Σ picks the branch guided by Q-structure (as computed in $\mathcal{M}_\infty$).*
- *If $\mathcal{T}$ is maximal then $\Sigma_\infty(\mathcal{T})$ = the unique cofinal branch b which moves $\tau_{A,0}^{\mathcal{M}_\infty}$ correctly for all $A \in OD$ such that there is some $(\mathcal{P}, \Sigma, A) \in \mathcal{F}$ i.e. for each such A, $i_b(\tau_{A,0}^{\mathcal{M}_\infty}) = \tau_{A^*,0}^{\mathcal{M}_b^{\mathcal{T}}}$.*

Lemma 3.21. *Given any such $\mathcal{T}$ as above, $\Sigma_\infty(\mathcal{T})$ exists.*

Proof. Suppose not. Again reflect the failure to a model N coded by a Suslin co-Suslin set. We may assume $N \vDash V = K^{\Sigma}(\mathbb{R})$ where $(\mathcal{P}, \Sigma)$ is a hod pair giving us $HOD|\theta_\alpha$. Just as in the previous proposition, we then get a next mouse $\mathcal{N}$ that iterates out to (possibly a longer mouse than) $\mathcal{M}_\infty^N$. This mouse $\mathcal{N}$ has strategy Λ with property that for all $A \in \mathbb{B}(P, \Sigma)$, there is a Λ-iterate $(\mathcal{M}, \Lambda_{\mathcal{M}})$ of $\mathcal{N}$ such that $\Lambda_{\mathcal{M}}$ strongly respects A. This easily gives us a contradiction. ☐

It is evident that $L(\mathcal{M}_\infty, \Sigma_\infty) \subseteq HOD$. Next, we show $\mathcal{M}_\infty$ and Σ_∞ capture all unbounded subsets of Θ in HOD. In $L(\mathcal{M}_\infty, \Sigma_\infty)$, first construct (using Σ_∞) a mouse $\mathcal{M}_\infty^+$ extending $\mathcal{M}_\infty$ such that $\mathrm{o}(\mathcal{M}_\infty)$ is the largest cardinal of $\mathcal{M}_\infty^+$ as follows:

1. Let $\mathbb{R}_G^*$ be the symmetric reals obtained from a generic G over $\mathcal{M}_\infty$ of $\mathrm{Col}(\omega, < \lambda^{\mathcal{M}_\infty})$.
2. For each A_G^* (defined as above) (we know $L(\mathbb{R}_G^*, A_G^*) \vDash AD^+$), pull back the hybrid mice over $\mathbb{R}_G^*$ in this model to hybrid mice $\mathcal{S}$ extending $\mathcal{M}_\infty$ with $D(S, \lambda^{M_\infty}) = L(\mathbb{R}_G^*, A_G^*)$.
3. Let $\mathcal{M}_\infty^+ = \cup_{\mathcal{S}} \mathcal{S}$ for all such $\mathcal{S}$ above. $\mathcal{M}_\infty^+$ is independent of G. By a reflection argument (and Prikry-like forcing) as above, the translated mice over $\mathcal{M}_\infty$ are all compatible, no levels of $\mathcal{M}_\infty^+$ projects across $o(\mathcal{M}_\infty)$, and $\mathcal{M}_\infty^+$ contains as its initial segments all translation of $\mathbb{R}_G^*$-mice in $D(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty})$. This is just saying that $\mathcal{M}_\infty^+$ contains enough mice to compute HOD.

Remark 3.22. Θ is not collapsed by Σ_∞ as it is a cardinal in HOD. Σ_∞ is used to obtain the A_G^* above by moving correctly the $\tau_{A,0}^{M_\infty}$ in genericity iterations. $L(\mathcal{M}_\infty)$ does not see the sequence $\langle \tau_{A,k}^{\mathcal{M}_\infty} | k \in \omega \rangle$ hence can't construct A_G^*. Also since Σ_∞ collapses $\delta_1^{M_\infty}, \delta_2^{\mathcal{M}_\infty}$..., it doesn't make sense to talk about $D(L(\mathcal{M}_\infty, \Sigma_\infty))$.

Lemma 3.23. *$HOD \subseteq L(\mathcal{M}_\infty, \Sigma_\infty)$*

Proof. Using Theorem 2.10, we know $HOD = L[P]$ for some $P \subseteq \Theta$. Therefore, it is enough to show $P \in L(\mathcal{M}_\infty, \Sigma_\infty)$. Let ϕ be a formula defining P, i.e.

$$\alpha \in P \Leftrightarrow V \vDash \phi[\alpha].$$

We suppress the ordinal parameter here. Now in $L(\mathcal{M}_\infty, \Sigma_\infty)$ let $\pi : \mathcal{M}_\infty|(\eta_0^{++})^{\mathcal{M}_\infty} \to (\mathcal{M}_\infty)^{(\mathcal{M}_\infty^+, \lambda^{\mathcal{M}_\infty})}$ where π is according to Σ_∞.

Claim: $\alpha \in P \Leftrightarrow D(\mathcal{M}_\infty^+, \lambda^{M_\infty}) \vDash \phi[\pi(\alpha)]$. $(*)$

Proof. Otherwise, reflect the failure of $(*)$ as before to get a model N coded by a Suslin co-Suslin set, a hod pair $(\mathcal{P}, \Sigma)$ giving us $HOD|\theta_\alpha$ such that

$$N \vDash ZF + DC + AD^+ + V = K^\Sigma(\mathbb{R}) + (\exists\alpha)(\phi[\alpha] \not\Leftrightarrow D(\mathcal{M}^+_\infty, \Sigma_\infty) \vDash \phi[\pi(\alpha)]).$$

Fix such an α. As before, let $\mathcal{N}$ be the next mouse (i.e. $\mathcal{N}$ has ω Woodins $\langle \delta_i \mid i < \omega \rangle$ on top of $\mathcal{P}$) with $\rho(\mathcal{N}) < sup_i \delta_i$) with strategy Λ extending Σ and Λ has branch condensation and is Ω-fullness preserving, where $\Omega = (\Sigma^2_1)^N$. We may assume Λ is guided by $\vec{D}$ where $\vec{D} = \langle D_n \mid n < \omega \rangle$ is defined as in Lemma 3.18. As before, we may assume $\mathcal{N}$ has derived model N. Let $\sigma : N|((\delta_0^N)^{++})^N \rightarrow (M_\infty)^{D(N,\lambda^N)}$ be the direct limit map by Λ. We may assume $\sigma(\overline{\alpha}) = \alpha$ for some $\overline{\alpha}$. Working in N, it then remains to see that:

$$D(\mathcal{M}^+_\infty, \lambda^{\mathcal{M}_\infty}) \vDash \phi[\pi(\alpha)] \Leftrightarrow D(N, \lambda^N) \vDash \phi[\sigma(\overline{\alpha})] \quad (**).$$

To see that $(**)$ holds, we need to see that the fragment of Λ that defines $\sigma(\overline{\alpha})$ can be defined in $D(N, \lambda^N)$. This then will give the equivalence in $(**)$. Because $\alpha < \eta_0$, $\overline{\alpha} < \delta_0$, pick an n such that such that $\gamma^{N,0}_{D_n,0} > \overline{\alpha}$. Then the fragment of Λ that defines $\sigma(\overline{\alpha})$ is definable from D_n (and $N|(\delta_0^N)$) in $D(N, \lambda^N)$, which is what we want.

The equivalence $(**)$ gives us a contradiction. □

The claim finishes the proof of $P \in L(\mathcal{M}_\infty, \Sigma_\infty)$. This then implies $HOD = L[P] \subseteq L(\mathcal{M}_\infty, \Sigma_\infty)$. □

Lemma 3.23 implies $HOD = L(\mathcal{M}_\infty, \Sigma_\infty)$, hence completes our computation.

4 The $\Theta = \theta_\alpha$, where α is limit, case

There are two cases: the easier case is when $HOD \vDash$ "$cof(\Theta)$ is not measurable", and the harder case is when $HOD \vDash$ "$cof(\Theta)$ is measurable".

Here's the direct limit system that gives us V^{HOD}_Θ.

$$\mathcal{F} = \{(\mathcal{Q}, \Lambda) \mid (\mathcal{Q}, \Lambda) \text{ is a hod pair and } \Lambda \text{ is fullness preserving and has branch condensation}\}.$$

The order on $\mathcal{F}$ is given by

$$(\mathcal{Q}, \Lambda) \leq^{\mathcal{F}} (\mathcal{R}, \Psi) \ iff \ \mathcal{Q} \text{ iterates to a hod initial segment of } \mathcal{R}.$$

By Theorem 3.4, $\leq^{\mathcal{F}}$ is directed and we can form the direct limit of $\mathcal{F}$ under the natural embeddings coming from the comparison process. Let $\mathcal{M}_\infty$ be the direct limit. By the computation in [3],

$$\mathcal{M}_\infty = V_\Theta^{HOD}.$$

We quote a theorem from [3] which will be used in the upcoming computation. For unexplained notations, see [3].

Theorem 4.1 (Sargsyan, Theorem 4.2.23 in [3])**.** *Suppose* $(\mathcal{P}, \Sigma)$ *is a hod pair such that* Σ *has branch condensation and is fullness preserving. There is then* $\mathcal{Q}$ *a* Σ*-iterate of* $\mathcal{P}$ *such that whenever* $\mathcal{R}$ *is a* $\Sigma_{\mathcal{Q}}$*-iterate of* $\mathcal{Q}$*,* $\alpha < \lambda^{\mathcal{R}}$*, and* $B \in (\mathbb{B}(\mathcal{R}(\alpha), \Sigma_{\mathcal{R}(\alpha)}))^{L(\Gamma(\mathcal{R}(\alpha+1), \Sigma_{\mathcal{R}(\alpha+1)}))}$

1. $\Sigma_{\mathcal{R}(\alpha+1)}$ *is super fullness preserving and is strongly guided by some*

$$\vec{B} = \langle B_i \mid i < \omega \rangle \subseteq (\mathbb{B}(\mathcal{R}(\alpha), \Sigma_{\mathcal{R}(\alpha)}))^{L(\Gamma(\mathcal{R}(\alpha+1), \Sigma_{\mathcal{R}(\alpha+1)}))};$$

2. *there is a* $\mathcal{S} \in pI(\mathcal{R}(\alpha+1), \Sigma_{\mathcal{R}(\alpha+1)})$ *such that* $\Sigma_{\mathcal{S}}$ *respects* B

We deal with the easy case first.

4.1 The nonmeasurable cofinality case

The following theorem is the full HOD computation in this case.

Theorem 4.2. $HOD = L(V_\Theta^{HOD})$

Proof. Technically, one needs the sequence of strategies of hod initial segments of $\mathcal{M}_\infty$ to compute the full HOD. The notation above however suggests that we don't need to explicitly add the sequence of strategies as an extra predicate as $\mathcal{M}_\infty$ can compute the sequence of strategies of its own hod initial segments.

To prove the theorem, suppose the equality is false. Then by Theorem 2.10, there is an $A \subseteq \Theta$ such that $A \in HOD \backslash L(V_\Theta^{HOD})$ (the fact that $L(V_\Theta^{HOD}) \subseteq HOD$ is obvious). By Σ_1-reflection (i.e. Theorem 2.8), there is a transitive N coded by a Suslin co-Suslin set such that

$$\begin{aligned} N \ \vDash \ & ZF^- + AD^+ V = L(\mathcal{P}(\mathbb{R})) + SMC + \text{“}\Theta \text{ exists and is limit in the Solovay sequence ”} + \\ & \text{“}HOD \vDash cof(\Theta) \text{ is not measurable ”} + \text{“}\exists B \subseteq \Theta (B \in HOD \backslash L(V_\Theta^{HOD}))\text{”}. \end{aligned}$$

Take N to be the minimal such and let B witness the failure of the theorem in N. Let ϕ

define B (for simplicity, we suppress the ordinal parameter) i.e.

$$\alpha \in B \; iff \; N \vDash \phi[\alpha]$$

Let $\Omega = \mathcal{P}(\mathbb{R})^N$. There is a pair $(\mathcal{P}, \Sigma)$ such that:

1. $\mathcal{P} = L_\beta(\cup_{\gamma<\lambda^P} P_\gamma)$ for some λ^P;
2. for all $\gamma < \lambda^{\mathcal{P}}$, $\mathcal{P}_\beta$ is a hod mouse whose strategy $\Sigma_\gamma \in \Omega$ is Ω-fullness preserving, has branch condensation, and $\lambda^{\mathcal{P}_\beta} = \beta$;
3. if $\gamma < \eta < \lambda^{\mathcal{P}}$, $\mathcal{P}_\gamma \trianglelefteq_{hod} \mathcal{P}_\eta$;
4. β is least such that $\rho_\omega(L_\beta(\cup_{\gamma<\lambda^{\mathcal{P}}}\mathcal{P}_\gamma)) < o(\cup_{\gamma<\lambda^{\mathcal{P}}}\mathcal{P}_\gamma))$;
5. $\mathcal{P} \vDash cof(\lambda^{\mathcal{P}})$ is not measurable;
6. Σ has branch condensation and extends $\oplus_{\gamma<\lambda^{\mathcal{P}}}\Sigma_\gamma$;

Such a $(\mathcal{P}, \Sigma)$ can be obtained by performing a Ω-hod pair construction (see Definition 3.10) inside some N_x^* capturing a good scaled pointclass beyond Ω. We may and do assume that $(\cup_{\gamma<\lambda^{\mathcal{P}}}\mathcal{P}_\gamma, \oplus_{\gamma<\lambda^{\mathcal{P}}}\Sigma_\gamma)$ satisfies Theorem 4.1 applied in N. This implies that the direct limit $\mathcal{M}_\infty^+$ of all Σ-iterates of $\mathcal{P}$ is a subset of HOD^N. Let $j : \mathcal{P} \to \mathcal{M}_\infty^+$ be the natural map. Then in N, $\mathcal{M}_\infty^+ | j(\lambda^{\mathcal{P}}) = V_\Theta^{HOD} = \mathcal{M}_\infty$.

Now pick a sequence $\langle \gamma_i \mid i < \omega \rangle$ cofinal in $\lambda^{\mathcal{P}}$ such that $\delta_{\lambda^{\mathcal{P}_{\gamma_i}}}$ is Woodin in $\mathcal{P}$, an enumeration $\langle x_i \mid i < \omega \rangle$ of $\mathbb{R}$ and do a genericity iteration of $\mathcal{P}$ to successively make each x_i generic at appropriate image of $\delta_{\lambda^{\mathcal{P}_{\gamma_i}}}$. Let $\mathcal{Q}$ be the end model of this process and $i : \mathcal{P} \to \mathcal{Q}$ be the iteration embedding. Then by assumption (5) above, we have that N is the derived model of $\mathcal{Q}$ at $i(\lambda^P)$.

In N, let D be the derived model of $\mathcal{M}_\infty^+$ at Θ and

$$\pi_\infty : \mathcal{M}_\infty \to (\mathcal{M}_\infty)^D$$

be the direct limit embedding given by the join of the strategies of $\mathcal{M}_\infty$'s hod initial segments. Then by the same argument as that given in Lemma 2.18, we have

$$\alpha \in B \; iff \; D \vDash \phi[\pi_\infty(\alpha)].$$

The proof of Lemma 2.18 also gives that $B \in (L(\mathcal{M}_\infty))^N$, which contradicts our assumption. Hence we're done. □

4.2 The measurable cofinality case

Suppose $HOD \vDash cof(\Theta)$ is measurable. We know by [3] that V_Θ^{HOD} is $\mathcal{N}_\infty$ where $\mathcal{N}_\infty$ is the direct limit (under the natural maps) of $\mathcal{F}$, where $\mathcal{F}$ is introduced at the beginning of this section. Let

$$\mathcal{M}_\infty = Ult_0(HOD, \mu)|\Theta,$$

where μ is the order zero measure on $cof^{HOD}(\Theta)$. For notational simplicity, for each $\beta < cof^{HOD}(\Theta) =_{def} \alpha$, let Λ_β be the strategy of $\mathcal{M}_\infty(\beta)$ and Σ_β be the strategy of $\mathcal{N}_\infty(\beta)$. Let

$$\mathcal{M}_\infty^+ = Ult_0(HOD, \mu)|(\Theta^+)^{Ult_0(HOD,\mu)},$$

and

$$\mathcal{N}_\infty^+ = \cup\{\mathcal{M} \mid \mathcal{N}_\infty \trianglelefteq \mathcal{M},\ \rho(\mathcal{M}) = \Theta, \mathcal{M} \text{ is a hybrid mouse satisfying property } (*)\}.$$

Here a mouse $\mathcal{M}$ satisfying property $(*)$ if whenever $\pi : \mathcal{M}^* \to \mathcal{M}$ is elementary and $\pi(\Theta^*) = \Theta$, then $\mathcal{M}^*$ is a $\oplus_{\alpha<\Theta^*}\Sigma^*_\alpha$-mouse for stacks above Θ^*, where $\Sigma^*_\alpha = \Sigma^\pi_{\pi(\alpha)}$ is the strategy for the hod mouse $\mathcal{M}^*(\alpha)$. We give two characterizations of HOD here: one in terms of $\mathcal{M}^+_\infty$ and the other in terms of $\mathcal{N}^+_\infty$. The first one is easier to see.

Theorem 4.3. *1. $HOD = L(V_\Theta^{HOD}, \mathcal{M}^+_\infty)$.*

2. $HOD = L(\mathcal{N}^+_\infty)$.

Proof. To prove (1), first let $j_\mu : HOD \to Ult_0(HOD, \mu)$ be the canonical ultrapower map. Let $A \in HOD$, $A \subseteq \Theta$. By the computation of HOD below Θ, we know that for each limit $\beta < \alpha$,

$$A \cap \theta_\beta \in Lp^{\oplus_{\gamma<\beta}\Sigma_\beta}(HOD|\theta_\beta).$$

From this, it is easy to see that there is a sequence $\langle \mathcal{M}_\beta \mid \beta < \alpha, \beta \text{ is limit}\rangle \in HOD$ such that for each limit $\beta < \alpha$

1. $\mathcal{M}_\beta$ is a $\oplus_{\gamma<\beta}\Sigma_\beta$-premouse over $HOD|\theta_\beta$;
2. $\mathcal{M}_\beta$ is countably iterable and $\rho(\mathcal{M}_\beta) = \theta_\beta$;
3. $A \cap \theta_\beta \in \mathcal{M}_\beta$.

Let $\mathcal{M}_\alpha = j_\mu(\langle \mathcal{M}_\beta \mid \beta < \alpha, \beta \text{ is limit}\rangle)(\alpha)$. Then $\mathcal{M}_\alpha \in \mathcal{M}^+_\infty$ and for any ordinal $\gamma < \Theta$

$$\gamma \in A \Leftrightarrow j_\mu(\gamma) \in \mathcal{M}_\alpha \Leftrightarrow (j_\mu|\Theta)(\gamma) \in \mathcal{M}_\alpha.$$

Since $j_\mu|\Theta$ agrees with the canonical ultrapower map $k : V_\Theta^{HOD} \to Ult_0(V^{HOD_\Theta}, \mu)$ on all ordinals less than Θ, the above equivalence shows that $A \in L(V_\Theta^{HOD}, \mathcal{M}_\infty^+)$. This proves (1).

The proof of (2) presented here is an elaboration of ideas communicated to the author by Grigor Sargsyan and John Steel. Suppose the statement of (2) is false. There is an $A \subseteq \Theta$ such that $A \in HOD \backslash \mathcal{N}_\infty^+$. By Σ_1-reflection (i.e. Theorem 2.8), there is a transitive N coded by a Suslin co-Suslin set such that

$$\begin{aligned} N \ \vDash \ & ZF^- + DC + V = L(\mathcal{P}(\mathbb{R})) + SMC + \text{“}\Theta \text{ exists and is limit in the Solovay sequence ”} \\ & + \text{“}HOD \vDash cof(\Theta) = \alpha \text{ is measurable ”} + \text{“}\exists A \subseteq \Theta(A \in HOD\backslash\mathcal{N}_\infty^+)\text{”}. \end{aligned}$$

Take N to be the minimal such and let A witness the failure of (2) in N. Let μ, j_μ, $\mathcal{M}_\infty$, $\mathcal{M}_\infty^+$, $\mathcal{N}_\infty$, $\mathcal{N}_\infty^+$ be as above but relativized to N. Working in N, as above there is a sequence $\langle \mathcal{M}_\beta \mid \beta < \alpha, \beta \text{ is limit}\rangle \in HOD$ such that for each limit $\beta < \alpha$

1. $\mathcal{M}_\beta$ is a $\oplus_{\gamma<\beta}\Sigma_\beta$-premouse over $HOD|\theta_\beta$;
2. $\mathcal{M}_\beta$ is countably iterable and $\rho(\mathcal{M}_\beta) = \theta_\beta$;
3. $A \cap \theta_\beta \in \mathcal{M}_\beta$.

Let $\Omega = \mathcal{P}(\mathbb{R})^N$. Fix an N_x^* capturing a good scaled pointclass beyond Ω. Now, we again do the Ω-hod pair construction in N_x^* to obtain a pair $(\mathcal{Q}, \Psi)$ such that

1. there is a limit ordinal $\lambda^\mathcal{Q}$ such that for all $\gamma < \lambda^\mathcal{Q}$, $\mathcal{Q}_\beta$ is a hod mouse with $\lambda^{\mathcal{Q}_\beta} = \beta$ and whose strategy $\Psi_\gamma \in \Omega$ is Ω-fullness preserving, has branch condensation;
2. if $\gamma < \eta < \lambda^\mathcal{Q}$, $\mathcal{Q}_\gamma \trianglelefteq_{hod} \mathcal{Q}_\eta$;
3. $\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma \vDash \mathrm{cof}(\lambda^\mathcal{Q})$ is measurable;
4. the induced strategy Λ of $Lp^{\Omega,\oplus_{\gamma<\lambda^\mathcal{Q}}\Psi_\gamma}(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)$[15] is not in Ω;
5. $\mathcal{Q}$ is a sound mouse from the $L[E, \oplus_{\gamma<\lambda^\mathcal{Q}}\Psi_\gamma][\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma]$-construction done in N_x^* that has projectum $\leq o(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)$; in fact, let $\mathcal{Q}$ be the first level in the construction with that property and Ψ be the induced strategy of $\mathcal{Q}$.

Let $\delta_{\lambda^\mathcal{Q}} = o(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)$ and $\eta = o(Lp^{\Omega,\oplus_{\gamma<\lambda^\mathcal{Q}}\Psi_\gamma}(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)) = (\delta_{\lambda^\mathcal{Q}}^+)^\mathcal{Q}$ and $\mathcal{M}_\infty(\mathcal{Q}, \Lambda)$ be the direct limit (under natural embeddings) of Λ-iterates of $\mathcal{Q}$.

Lemma 4.4. *$\mathcal{M}_\infty(Q, \Lambda)$ exists.*

---

[15] If $\mathcal{M} \lhd Lp^{\Omega,\oplus_{\gamma<\lambda^\mathcal{Q}}\Psi_\gamma}(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)$ and $\mathcal{M}$ extends $\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma$ then $\mathcal{M}$ is a mouse in N in the sense that N knows how to iterate $\mathcal{M}$ for stacks above $o(\cup_{\gamma<\lambda^\mathcal{Q}}\mathcal{Q}_\gamma)$.

Proof. First note that Λ is Ω-fullness preserving. To see this, suppose not. Let $k : \mathcal{Q} \to \mathcal{R}$ be according to Λ witnessing this. It's easy to see that the tail $\Lambda_{\mathcal{R}}$ of Λ acting on $\mathcal{R}|k(\eta)$ is not in Ω (otherwise, $\Lambda^k_{\mathcal{R}} = \Lambda$ by hull condensation and hence $\Lambda \in \Omega$. Contradiction.) However, $\oplus_{\gamma<\lambda^{\mathcal{R}}}\Psi_{\mathcal{R}(\gamma)} \in \Omega$ since the iterate of N^*_x by the lift-up of k thinks that the fragment of its strategy inducing $\oplus_{\gamma<\lambda^{\mathcal{R}}}\Psi_{\mathcal{R}(\gamma)}$ is in Ω. Now suppose $\mathcal{M}$ is a $\oplus_{\gamma<\lambda^{\mathcal{R}}}\Psi_{\mathcal{R}(\gamma)}$-mouse projecting to $\delta_{\lambda^{\mathcal{R}}}$ with strategy Ξ in Ω and $\mathcal{M} \ntrianglelefteq \mathcal{R}$ (again, Ξ acts on trees above $\delta_{\lambda^{\mathcal{R}}}$ and moves the predicates for $\oplus_{\gamma<\lambda^{\mathcal{R}}}\Psi_{\mathcal{R}(\gamma)}$ correctly). We can compare $\mathcal{M}$ and $\mathcal{R}$ (the comparison is above $\delta_{\lambda^{\mathcal{R}}}$). Let $\overline{\mathcal{M}}$ be the last model on the $\mathcal{M}$ side and $\overline{\mathcal{R}}$ on the $\mathcal{R}$ side. Then $\overline{\mathcal{R}} \lhd \overline{\mathcal{M}}$. Let $\pi : \mathcal{R} \to \overline{\mathcal{R}}$ be the iteration map from the comparison process and Σ be the $\pi \circ k$-pullback of the strategy of $\overline{\mathcal{R}}$. Hence $\Sigma \in \Omega$ since $\Xi \in \Omega$. Σ acts on trees above $\delta_{\lambda^{\mathcal{Q}}}$ and moves the predicate for $\oplus_{\gamma<\lambda^{\mathcal{Q}}}\Psi_\gamma$ correctly by by our assumption on Ξ and branch condensation of $\oplus_{\gamma<\lambda^{\mathcal{Q}}}\Psi_\gamma$. These properties of Σ imply that $\mathcal{Q} \lhd Lp^{\Omega,\oplus_{\gamma<\lambda^{\mathcal{Q}}}\Psi_\gamma}(\cup_{\gamma<\lambda^{\mathcal{Q}}}\mathcal{Q}_\gamma)$. Contradiction. For the case that there are $\alpha < \lambda^R$, $\delta^R_\alpha \leq \eta < \delta^R_{\eta+1}$, and η is a strong cutpoint of R, and $\mathcal{M}$ is a sound $\Psi_{R(\alpha)}$-mouse projecting to η with iteration strategy in Ω, the proof is the same as that of Theorem 3.7.6 in [3].

Now we show Λ has branch condensation. The proof of this is due to John Steel. For notational simplicity, we write Λ^- for $\oplus_{\gamma<\lambda^{\mathcal{Q}}}\Psi_\gamma$. Hence, $\Lambda \notin \Omega$ and $\Lambda^- \in \Omega$. Suppose Λ does not have branch condensation. We have a minimal counterexample as follows: there are an iteration $i : \mathcal{Q} \to \mathcal{R}$ by Λ, a normal tree $\mathcal{U}$ on $\mathcal{R}$ in the window $[\xi, \gamma)$ where $\xi < \gamma$ are two consecutive Woodins in $\mathcal{R}$ such that $\sup i''\delta_{\lambda^{\mathcal{Q}}} \leq \xi$, two distinct cofinal branches of $\mathcal{U}$: b and $c = \Lambda_{\mathcal{R}}(\mathcal{U})$, an iteration map $j : \mathcal{Q} \to \mathcal{S}$ by Λ, and a map $\sigma : \mathcal{M}^{\mathcal{U}}_b \to \mathcal{S}$ such that $j = \sigma \circ i^{\mathcal{U}}_b \circ i$. We may also assume that if $\overline{\mathcal{R}}$ is a model along the main branch of the stack from $\mathcal{Q}$ to $\mathcal{R}$ giving rise to i and $i_{\overline{\mathcal{R}},\mathcal{R}} : \overline{\mathcal{R}} \to \mathcal{R}$ be the natural map such that $i_{\overline{\mathcal{R}},\mathcal{R}}(\overline{\xi}) = \xi$ and $i_{\overline{\mathcal{R}},\mathcal{R}}(\overline{\gamma}) = \gamma$ then the extenders used to get from $\mathcal{Q}$ to $\overline{\mathcal{R}}$ have generators below $\overline{\xi}$. This gives us $\sup(Hull^{\mathcal{R}}(\xi \cup \{p\}) \cap \gamma) = \gamma$ where p is the standard parameter of $\mathcal{R}$. Let $\Phi = \Lambda^\sigma_{\mathcal{S}}$ and Φ^- have the obvious meaning. It's easy to see that $\Phi \notin \Omega$ and $\Phi^- \in \Omega$. By the same proof as in the previous paragraph, Φ is Ω-fullness preserving. This of course implies that $\mathcal{M}^{\mathcal{U}}_b$ is Ω-full. Now we compare $\mathcal{M}^{\mathcal{U}}_b$ and $\mathcal{M}^{\mathcal{U}}_c$. First we line up the strategies of $\mathcal{M}^{\mathcal{U}}_b|\delta(\mathcal{U})$ and $\mathcal{M}^{\mathcal{U}}_c|\delta(\mathcal{U})$ by iterating them into the (Ω-full) hod pair construction of some N^*_y (where y codes $(x, \mathcal{M}^{\mathcal{U}}_c, \mathcal{M}^{\mathcal{U}}_b)$). This process produces a single normal tree $\mathcal{W}$. Let $a = \Phi(\mathcal{W})$ and $d = \Lambda_{\mathcal{M}^{\mathcal{U}}_c}(\mathcal{W})$. Let $X = Hull^{\mathcal{R}}(\xi \cup \{p\}) \cap \gamma$. Note that $(i^{\mathcal{W}}_a \circ i^{\mathcal{U}}_b)\text{''}\mathrm{X} \subseteq \delta(\mathcal{W})$ and $i^{\mathcal{W}}_d \circ i^{\mathcal{U}}_c\text{''}\mathrm{X} \subseteq \delta(\mathcal{W})$. Now continue lining up $\mathcal{M}^{\mathcal{W}}_a$ and $\mathcal{M}^{\mathcal{W}}_d$ above $\delta(\mathcal{W})$ (using the same process as above). We get $\pi : \mathcal{M}^{\mathcal{W}}_a \to \mathcal{K}$ and $\tau : \mathcal{M}^{\mathcal{W}}_d \to \mathcal{K}$ (we indeed end up with the same model $\mathcal{K}$ by our assumption on the pair (Λ, Λ^-)). But then

$$(\pi \circ i^{\mathcal{W}}_a \circ i^{\mathcal{U}}_b)\text{''}\mathrm{X} = (\tau \circ i^{\mathcal{W}}_d \circ i^{\mathcal{U}}_c)\text{''}\mathrm{X}.$$

But by the fact that $(i^{\mathcal{W}}_a \circ i^{\mathcal{U}}_b)$"X $\subseteq \delta(\mathcal{W})$ and $i^{\mathcal{W}}_d \circ i^{\mathcal{U}}_c$"X $\subseteq \delta(\mathcal{W})$ and π agrees with τ above $\delta(\mathcal{W})$, we get

$$(i^{\mathcal{W}}_a \circ i^{\mathcal{U}}_b)\text{"X} = (i^{\mathcal{W}}_d \circ i^{\mathcal{U}}_c)\text{"X}.$$

This gives $ran(i^{\mathcal{W}}_a) \cap ran(i^{\mathcal{W}}_d)$ is cofinal in $\delta(\mathcal{W})$, which implies $a = d$. This in turns easily implies $b = c$. Contradiction.

Finally, let $\mathcal{R}$ and $\mathcal{S}$ be Λ-iterates of $\mathcal{Q}$ and let $\Lambda_{\mathcal{R}}$ and $\Lambda_{\mathcal{S}}$ be the tails of Λ on $\mathcal{R}$ and $\mathcal{S}$ respectively. We want to show that $\mathcal{R}$ and $\mathcal{S}$ can be further iterated (using $\Lambda_{\mathcal{R}}$ and $\Lambda_{\mathcal{S}}$ respectively) to the same model. To see this, we compare $\mathcal{R}$ and $\mathcal{S}$ against the Ω-full hod pair construction of some N^*_y (for some y coding $(x, \mathcal{R}, \mathcal{S})$). Then during the comparison, only $\mathcal{R}$ and $\mathcal{S}$ move (to say $\mathcal{R}^*$ and $\mathcal{S}^*$). It's easy to see that $\mathcal{R}^* = \mathcal{S}^*$ and their strategies are the same (as the induced strategy of N^*_y on its appropriate background construction). □

By the properties of $(\mathcal{Q}, \Psi)$ and Λ, we get that $\rho(\mathcal{M}_\infty(\mathcal{Q}, \Lambda)) \leq \Theta$ and $(HOD|\Theta)^N = \mathcal{M}_\infty(\mathcal{Q}, \Lambda)|\Theta$. Let k be the least such that $\rho_{k+1}(\mathcal{Q}) \leq \delta_{\lambda^{\mathcal{Q}}}$.

Claim. $\mathcal{M}_\infty(\mathcal{Q}, \Lambda) \notin N$

Proof. Suppose not. Let $i : \mathcal{Q} \to \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$ be the direct limit map according to Λ. By an absoluteness argument (i.e. using the absoluteness of the illfoundedness of the tree built in $N[g]$ for $g \subseteq Col(\omega, |\mathcal{M}_\infty(\mathcal{Q}, \Lambda)|)$ generic over N of approximations of a embedding from $\mathcal{Q}$ into $\mathcal{M}_\infty(\mathcal{Q}, \Xi)$ extending the iteration embedding according to $\oplus_{\beta<\lambda^{\mathcal{Q}}}\Psi_\beta$ on $\mathcal{Q}|\delta_{\lambda^{\mathcal{Q}}}$), we get a map π such that

1. $\pi \in N$
2. $\pi : \mathcal{Q} \to \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$;
3. for each $\beta < \lambda^{\mathcal{Q}}$, $\pi|Q(\beta)$ is according to Ψ_β.
4. $\pi(p) = i(p)$ where $p = p_k(\mathcal{Q})$.

This implies that $\pi = i \in N$ since $\mathcal{Q}$ is sound and $\rho(\mathcal{Q}) \leq \delta^{\lambda^{\mathcal{Q}}}$. But this map determines Λ by branch condensation of Λ. So $\Lambda \in N$. But this is a contradiction. □

Returning to the proof of (2), let $j =_{def} j_\mu : HOD \to Ult_0(HOD, \mu)$ and $\mathcal{M}_\alpha = j(\langle \mathcal{M}_\beta \mid \beta < \alpha, \beta \text{ is limit}\rangle)(\alpha)$. Let $i : \mathcal{M}_\infty(\mathcal{Q}, \Lambda) \to Ult_k(\mathcal{M}_\infty(\mathcal{Q}, \Lambda), \mu)$ be the canonical map. Note that $A \notin \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$. To see this, assume not, let $\mathcal{R} \lhd \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$ be the first level $\mathcal{S}$ of $\mathcal{M}_\infty(\mathcal{Q}, \Lambda)$ such that A is definable over $\mathcal{S}$. Note that $\mathcal{R} \in N$. To get a contradiction, we show $\mathcal{R} \lhd \mathcal{N}^+_\infty$ by showing $\mathcal{R}$ is satisfies property $(*)$ in N. Let $\mathcal{K}$ be a countable mouse embeddable into $\mathcal{R}$ by a map $k \in N$. Then we can compare $\mathcal{K}$ and $\mathcal{Q}$ against the Ω-full hod pair construction of some N_y^* just like in the argument on the previous page; hence we may assume $\mathcal{K} \lhd \mathcal{Q}$ ($\mathcal{Q} \unlhd \mathcal{K}$ can't happen because then $\Lambda \in N$). The minimality assumption on Q easily implies $\mathcal{K} \lhd Lp^{\Omega,\oplus_{\gamma<\lambda^{\mathcal{Q}}}\Psi_\gamma}(\mathcal{Q}|\delta_{\lambda^{\mathcal{Q}}})$. But then N can iterate $\mathcal{K}$ for stacks on $\mathcal{K}$ above $\delta_{\lambda^{\mathcal{Q}}} = \delta_{\lambda^{\mathcal{K}}}$, which is what we want to show. The fact that $\mathcal{R} \lhd \mathcal{N}_\infty^+$ contradicts $A \notin \mathcal{N}_\infty^+$.

Next, we note that $Ult_0(HOD, \mu)|\Theta = Ult_k(\mathcal{M}_\infty(\mathcal{Q}, \Lambda), \mu)|\Theta$ and $i|\Theta = j|\Theta$. Let $\mathcal{R} = Th^{\mathcal{M}_\infty(\mathcal{Q},\Lambda)}(\Theta \cup \{p\})$ where $p = p_k(\mathcal{M}_\infty(\mathcal{Q}, \Lambda))$ and $\mathcal{S} = Th^{Ult_k(\mathcal{M}_\infty(\mathcal{Q},\Lambda),\mu)}(\Theta \cup \{i(p)\})$. We have that $\mathcal{M}_\alpha$ and $\mathcal{S}$ are sound hybrid mice in the same hierarchy, hence by countable iterability, we can conclude either $\mathcal{M}_\alpha \lhd \mathcal{S}$ or $\mathcal{S} \unlhd \mathcal{M}_\alpha$.

If $\mathcal{M}_\alpha \lhd \mathcal{S}$, then $\mathcal{M}_\alpha \in Ult_k(\mathcal{M}_\infty(\mathcal{Q}, \Lambda), \mu)$. This implies $A \in \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$ by a computation similar to that in the proof of (1), i.e.

$$\beta \in A \Leftrightarrow \mathcal{M}_\infty(\mathcal{Q}, \Lambda) \vDash (i|\Theta)(\beta) \in \mathcal{M}_\alpha.$$

This is a contradiction to the fact that $A \notin \mathcal{M}_\infty(\mathcal{Q}, \Lambda)$. Now suppose $\mathcal{S} \unlhd \mathcal{M}_\alpha$. This then implies $\mathcal{S} \in Ult_0(HOD, \mu)$, which in turns implies $\mathcal{M}_\infty(\mathcal{Q}, \Lambda) \in HOD$ by the following computation: for any formula ϕ and $s \in \Theta^{<\omega}$,

$$(\phi, s) \in \mathcal{R} \Leftrightarrow HOD \vDash (\phi, (j|\Theta)(s)) \in \mathcal{S}.$$

This is a contradiction to the claim. This completes the proof of (2). $\square$

Theorem 4.3 completes our analysis of HOD for natural determinacy models below "$AD_\mathbb{R} + \Theta$ is regular." Our analysis shows that essentially, the HOD's of class models satisfying "$AD^+ + V = L(\mathcal{P}(\mathbb{R}))$" form a coherent hierarchy of mice in that HOD of a such a model extends (as a mouse) those of smaller ones.

5 Questions and open problems

The following question is natural.

Question. Assume $AD^+ + V = L(\mathcal{P}_\mathbb{R})$. Does HOD satisfy GCH?

More generally (and vaguely), we can ask whether HOD is a fine-structural model. As shown in [3] and in this paper, under $(*)$, HOD is indeed a fine-structural model. The next natural determinacy theory to aim to understand HOD for seems to be the theory "$AD^+ + \Theta = \theta_{\alpha+1} + \theta_\alpha$ is the largest Suslin cardinal." It's not known whether this theory is consistent (relative to large cardinals). Recent (unpublished) work suggests that this theory is consistent assuming the existence of a mouse with a measurable Woodin cardinal. The best result to date on iterability is in [2] and falls just short of this.

References

[1] Steve Jackson. Structural consenquences of AD,. *Handbook of Set Theory.*

[2] Itay Neeman. Inner models in the region of a Woodin limit of Woodin cardinals. *Ann. Pure Appl. Logic*, 116(1-3):67–155, 2002.

[3] Grigor Sargsyan. A tale of hybrid mice, available at http://math.ucla.edu/∼grigor/.

[4] Ralf Schindler and John R. Steel. *The core model induction.* Available at math.berkeley.edu/∼steel.

[5] J. R. Steel. Scales in $K(\mathbb{R})$ at the end of a weak gap. *J. Symbolic Logic*, 73(2):369–390, 2008.

[6] John R. Steel. An outline of nner model theory, available at www.math.berkeley.edu/∼ steel, to appear in the *Hanbook of Set theory.*

[7] John R. Steel. A theorem of Woodin on mouse sets, available at http://math.berkeley.edu/ steel/.

[8] John R. Steel. Woodin's analysis of $\mathrm{HOD}^{\mathrm{L}(\mathbb{R})}$*, available at www.math.berkeley.edu/∼ steel.*

[9] John R. Steel. Scales in $K(\mathbb{R})$*.* The Cabal Seminar*, 1:176–208, 2003.*

[10] John R. Steel. Derived models associated to mice. In Computational prospects of infinity. Part I. Tutorials*, volume 14 of* Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 105–193. World Sci. Publ., Hackensack, NJ, 2008.*

[11] John R. Steel. An optimal consistency strength lower bound for $AD_{\mathbb{R}}$*. Unpublished notes, 2008.*

[12] W. Hugh Woodin. The axiom of determinacy, forcing axioms, and the nonstationary ideal*, volume 1 of* de Gruyter Series in Logic and its Applications*. Walter de Gruyter & Co., Berlin, 1999.*